

## Immunotag™ TNF-IP 8 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT4685
Product Description	Immunotag™ TNF-IP 8 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	TNFIP 8
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IHC-p,IF,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human,Mouse
Host Species	Rabbit
Immunogen	The antiserum was produced against synthesized peptide derived from human TFIP8. AA range:31-80
Specificity	TNF-IP 8 Polyclonal Antibody detects endogenous levels of TNF-IP 8 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	TNFAIP8
Accession No.	O95379 Q921Z5
Alternate Names	TNFAIP8; Tumor necrosis factor alpha-induced protein 8; TNF alpha-induced protein 8; Head and neck tumor and metastasis-related protein; MDC-3.13; NF-kappa-B-inducible DED-containing protein; NDED; SCC-S2; TNF-induced protein GG2-1

## Antibody Specification

Description	developmental stage:Expressed at high levels in the fetal liver, lung and kidney.,function:Acts as a negative mediator of apoptosis and may play a role in tumor progression. Suppresses the TNF-mediated apoptosis by inhibiting caspase-8 activity but not the processing of procaspase-8, subsequently resulting in inhibition of BID cleavage and caspase-3 activation.,induction:By nuclear factor-KB (NF-KB) and TNF. Induction by TNF depends upon activation of NF-KB.,similarity:Belongs to the TNFAIP8 family.,tissue specificity:Expressed at high levels in the spleen, lymph node, thymus, thyroid, bone marrow and placenta. Expressed at high levels both in various tumor tissues, unstimulated and cytokine-activated cultured cells. Expressed at low levels in the spinal cord, ovary, lung, adrenal glands, heart, brain, testis and skeletal muscle.,
Protein Expression	Brain,Heart,Placenta,Umbilical vein,Urinary bladder,
Subcellular Localization	cytoplasm,
Protein Function	developmental stage:Expressed at high levels in the fetal liver, lung and kidney.,function:Acts as a negative mediator of apoptosis and may play a role in tumor progression. Suppresses the TNF-mediated apoptosis by inhibiting caspase-8 activity but not the processing of procaspase-8, subsequently resulting in inhibition of BID cleavage and caspase-3 activation.,induction:By nuclear factor-KB (NF-KB) and TNF. Induction by TNF depends upon activation of NF-KB.,similarity:Belongs to the TNFAIP8 family.,tissue specificity:Expressed at high levels in the spleen, lymph node, thymus, thyroid, bone marrow and placenta. Expressed at high levels both in various tumor tissues, unstimulated and cytokine-activated cultured cells. Expressed at low levels in the spinal cord, ovary, lung, adrenal glands, heart, brain, testis and skeletal muscle.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.