

Immunotag™ TRAF2 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT4717
Product Description	Immunotag™ TRAF2 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	TRAF2
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IHC-p,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human
Host Species	Rabbit
Immunogen	Synthesized peptide derived from the Internal region of human TRAF2.
Specificity	TRAF2 Polyclonal Antibody detects endogenous levels of TRAF2 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	TRAF2
Accession No.	Q12933 P39429
Alternate Names	TRAF2; TRAP3; TNF receptor-associated factor 2; E3 ubiquitin-protein ligase TRAF2; Tumor necrosis factor type 2 receptor-associated protein 3

Antibody Specification

Description	TNF receptor associated factor 2(TRAF2) Homo sapiens The protein encoded by this gene is a member the TNF receptor associated factor (TRAF) protein family. TRAF proteins associate with, and mediate th signal transduction from members of the TNF receptor superfamily. This protein directly interacts with TNF receptors, and forms a heterodimeric complex with TRAF1. This protein is required for TNF-alpha-mediated activation of MAPK8/JNK and NF-kappaB. The protein complex formed by this protein and TRAF1 interacts with the inhibitor-of-apoptosis proteins (IAPs), and functions as a mediator of the anti-apoptotic signals from TNF receptors. The interaction of this protein with TRADD, a TNF receptor associated apoptotic signal transducer, ensures the recruitment of IAPs for the direct inhibition of caspase activation. BIRC2/c-IAP1, an apoptosis inhibitor possessing ubiquitin ligase activity, can ubiquitinate and induce the degradation of this pro
Cell Pathway/ Category	MAPK_ERK_Growth,MAPK_G_Protein,Apoptosis_Inhibition,Apoptosis_Mitochondrial,Apoptosis_Overview,I-like receptor,Adipocytokine,Pathways in cancer,Small cell lung cancer,
Protein Expression	Brain,Cerebellum,Colon,Fetal brain,Human endometrium,Kidney,Leukocyte,Splee
Subcellular Localization	ubiquitin ligase complex,intracellular,cytoplasm,cytosol,cell cortex,cytoplasmic side of plasma membrane,vesicle membrane,CD40 receptor complex,membrane raft,TRAF2-GSTP1 complex,AIP1-IRE1 complex,IRE1-TRAF2-ASK1 complex,
Protein Function	domain:The coiled coil domain mediates homo- and hetero-oligomerization.,domain:The MATH/TRAF domain binds to receptor cytoplasmic domains.,function:Adapter protein and signal transducer that link members of the tumor necrosis factor receptor family to different signaling pathways by association with the receptor cytoplasmic domain and kinases. Association to the receptor is also mediated by the interaction with TRADD. Mediates activation of NF-kappa-B and JNK and is involved in apoptosis. The TRAF1/TRAF2 complex recruits the apoptotic suppressors BIRC2 and BIRC3 to TNFRSF1B/TNFR2. Seems to be involved in IL-15 signaling.,PTM:Ubiquitinated; mediated by SIAH2 and leading to its subsequent proteasomal degradation. Not ubiquitinated by SIAH1.,similarity:Contains 1 MATH domain.,similarity:Contains 1 RING-type zinc finger.,similarity:Contains 2 TRAF-type zinc fingers.,subunit:Homotrimer (Probable). Heteromer with TRAF1. Binds to TNFRSF1B/TNFR2, TNFRSF4 and TNFRSF5/CD40. Associates with CD27, TNFRSF8/CD30, TNFRSF9/CD137, TNFRSF11A/RANK, TNFRSF13B/TACI. TNFRSF14, TNFRSF16/NGFR, TNFRSF17/BCMA, TNFRSF18/AITR, TNFRSF19/TROY, TNFRSF19L/RELT, XEDAR, EDAR, Epstein-Barr virus BNFL1/LMP-1, IL15RA, TANK/ITRAF, RIPK2, TNIK, MAP3K14, MAP3K5, MAP3K1, MAP4K2, CDK9, CSK, and TRAF-interacting protein TRAIIP and TRAF and TNF receptor associated protein TTRAP. Interacts with TNFAIP3 and TRPC4AP. Interacts with PEG3 (By similarity). Binds to TRADD. Interacts with BIRC2 and BIRC3 N-terminus. Interacts with CYLD and TBK1. Interacts with MAVS/IPS1. Interacts with CASP8AP2 and USP48. Interacts with DAB2IP. Interacts with NFATC2IP and with HIVP3.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.