

Immunotag™ Trypsin-3 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT4754
Product Description	Immunotag™ Trypsin-3 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	Trypsin-3
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human
Host Species	Rabbit
Immunogen	The antiserum was produced against synthesized peptide derived from human Trypsin-3. AA range:117-166
Specificity	Trypsin-3 Polyclonal Antibody detects endogenous levels of Trypsin-3 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	PRSS3
Accession No.	P35030
Alternate Names	PRSS3; PRSS4; TRY3; TRY4; Trypsin-3; Brain trypsinogen; Mesotrypsinogen; Serine protease 3; Serine protease 4; Trypsin III; Trypsin IV

Antibody Specification

Description	protease, serine 3(PRSS3) Homo sapiens This gene encodes a trypsinogen, which is a member of the trypsin family of serine proteases. This enzyme is expressed in the brain and pancreas and is resistant to common trypsin inhibitors. It is active on peptide linkages involving the carboxyl group of lysine or arginine. This gene is localized to the locus of T cell receptor beta variable orphans on chromosome 9. Four transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Oct 2010],
Cell Pathway/ Category	Neuroactive ligand-receptor interaction,
Protein Expression	Brain,Pancreas,PCR rescued clones,Prostate,
Subcellular Localization	extracellular region,extracellular space,extracellular exosome,
Protein Function	catalytic activity:Preferential cleavage: Arg- -Xaa, Lys- -Xaa.,cofactor:Binds 1 calcium ion per subunit.,similarity:Belongs to the peptidase S1 family.,similarity:Contains 1 peptidase S1 domain.,tissue specificity:Pancreas and brain.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.