

Immunotag™ UBTD2 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT4811
Product Description	Immunotag™ UBTD2 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	UBTD2
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human,Mouse
Host Species	Rabbit
Immunogen	The antiserum was produced against synthesized peptide derived from human UBTD2. AA range:181-230
Specificity	UBTD2 Polyclonal Antibody detects endogenous levels of UBTD2 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	UBTD2
Accession No.	Q8WUN7 Q6PGH0
Alternate Names	UBTD2; DCUBP; SB72; Ubiquitin domain-containing protein 2; Dendritic cell-derived ubiquitin-like protein; DC-UbP; Ubiquitin-like protein SB72
Description	similarity:Contains 1 ubiquitin-like domain.,tissue specificity:Detected in dendritic cells. Highly expressed in tumor cell lines, but not detectable in most tissues.,
Protein Expression	Dendritic cell,Mammary gland,

Antibody Specification

Subcellular Localization	cytoplasm,
Protein Function	similarity:Contains 1 ubiquitin-like domain.,tissue specificity:Detected in dendritic cells. Highly expressed in tumor cell lines, but not detectable in most tissues.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.