

Immunotag™ β-1,4-Gal-T5 Polyclonal Antibody

| Antibody Specification | |
|------------------------|--|
| Catalog No. | ITT5010 |
| Product Description | Immunotag™ β-1,4-Gal-T5 Polyclonal Antibody |
| Size | 50 µg, 100 µg |
| Conjugation | HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647 |
| IMPORTANT NOTE | This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return. |
| Target Protein | β-1,4-Gal-T5 |
| Clonality | Polyclonal |
| Storage/Stability | -20°C/1 year |
| Application | WB,ELISA |
| Recommended Dilution | Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. |
| Concentration | 1 mg/ml |
| Reactive Species | Human,Mouse |
| Host Species | Rabbit |
| Immunogen | The antiserum was produced against synthesized peptide derived from human B4GALT5. AA range:321-370 |
| Specificity | β-1,4-Gal-T5 Polyclonal Antibody detects endogenous levels of β-1,4-Gal-T5 protein. |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen |
| Form | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Gene Name | B4GALT5 |
| Accession No. | O43286 Q9JMK0 |
| Alternate Names | B4GALT5; Beta-1; 4-galactosyltransferase 5; Beta-1,4-GalTase 5; Beta4Gal-T5; b4Gal-T5; Beta-1,4-GalT II; UDP-Gal:beta-GlcNAc beta-1,4-galactosyltransferase 5; UDP-galactose:beta-N-acetylglucosamine beta-1,4-galactosyltransferase 5 |

Antibody Specification

| | |
|-----------------------------|--|
| Description | beta-1,4-galactosyltransferase 5(B4GALT5) Homo sapiens This gene is one of seven beta-1,4-galactosyltransferase (beta4GalT) genes. They encode type II membrane-bound glycoproteins that appear to have exclusive specificity for the donor substrate UDP-galactose; all transfer galactose in a beta1,4 linkage to similar acceptor sugars: GlcNAc, Glc, and Xyl. Each beta4GalT has a distinct function in the biosynthesis of different glycoconjugates and saccharide structures. As type II membrane proteins, they have an N-terminal hydrophobic signal sequence that directs the protein to the Golgi apparatus and which then remains uncleaved to function as a transmembrane anchor. By sequence similarity, the beta4GalTs form four groups: beta4GalT1 and beta4GalT2, beta4GalT3 and beta4GalT4, beta4GalT5 and beta4GalT6, and beta4GalT7. The function of the enzyme encoded by this gene is not clear. This gene was previously designated as B4GALT4 but was renamed |
| Cell Pathway/ Category | O-Glycan biosynthesis, |
| Protein Expression | Cerebellum,Lung,Mammary tumor,Placenta, |
| Subcellular Localization | Golgi membrane,integral component of membrane,Golgi cisterna membrane,extracellular exosome, |
| Protein Function | cofactor:Manganese.,function:Responsible for the synthesis of complex-type N-linked oligosaccharides in many glycoproteins as well as the carbohydrate moieties of glycolipids.,online information:Beta-1,4-galactosyltransferase 5,online information:GlycoGene database,pathway:Protein modification; protein glycosylation.,similarity:Belongs to the glycosyltransferase 7 family.,subcellular location:Trans cisternae of Golgi stack.,tissue specificity:Ubiquitously expressed., |
| Usage | For Research Use Only! Not for diagnostic or therapeutic procedures. |