

Immunotag™ PREP Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT5132
Product Description	Immunotag™ PREP Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	PREP
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IHC-p,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human,Mouse,Rat
Host Species	Rabbit
Immunogen	Synthesized peptide derived from PREP . at AA range: 10-90
Specificity	PREP Polyclonal Antibody detects endogenous levels of PREP protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	PREP
Accession No.	Q8N6D4
Alternate Names	PREP; PEP; Prolyl endopeptidase; PE; Post-proline cleaving enzyme
Description	prolyl endopeptidase(PREP) Homo sapiens The protein encoded by this gene is a cytosolic prolyl endopeptidase that cleaves peptide bonds on the C-terminal side of prolyl residues within peptides that are up to approximately 30 amino acids long. Prolyl endopeptidases have been reported to be involved in the maturation and degradation of peptide hormones and neuropeptides. [provided by RefSeq, Jul 2008],

Antibody Specification

Protein Expression	Blood,Brain,Lymphocyte,Skin,
Subcellular Localization	nucleus,cytoplasm,membrane,
Protein Function	catalytic activity:Hydrolysis of Pro-[Xaa >> Ala-[Xaa in oligopeptides.,function:Cleaves peptide bonds on the C-terminal side of prolyl residues within peptides that are up to approximately 30 amino acids long.,PTM:The N-terminus is blocked.,similarity:Belongs to the peptidase S9A family.,subunit:Monomer.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.