

Immunotag™ VGF Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT5296
Product Description	Immunotag™ VGF Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	VGF
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IHC-p,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. IHC-p: 1/100-1/300. ELISA: 1/20000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human,Mouse,Rat
Host Species	Rabbit
Immunogen	Synthesized peptide derived from Neurosecretory protein VGF at AA range: 441-490
Specificity	VGF Polyclonal Antibody detects endogenous levels of VGF protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	VGF
Accession No.	O15240 P20156
Alternate Names	VGF; Neurosecretory protein VGF

Antibody Specification

Description	VGF nerve growth factor inducible(VGF) Homo sapiens This gene is specifically expressed in a subpopulation of neuroendocrine cells, and is upregulated by nerve growth factor. The structural organization of this gene is similar to that of the rat gene, and both the translated and the untranslated regions show a high degree of sequence similarity to the rat gene. The encoded secretory protein also shares similarities with the secretogranin/chromogranin family, however, its exact function is not known. [provided by RefSeq, Jul 2008],
Protein Expression	Brain,Placenta,PNS,
Subcellular Localization	extracellular space,transport vesicle,cytoplasmic vesicle,intracellular membrane-bounded organelle,
Protein Function	function:May be involved in the regulation of cell-cell interactions or in synatogenesis during the maturation of the nervous system.,subcellular location:Stored in secretory vesicles and then secreted.,tissue specificity:Expressed in brain.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.