

Immunotag™ OAS2 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT5536
Product Description	Immunotag™ OAS2 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	OAS2
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IHC-p,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. IHC-p: 1:100-1:300. ELISA: 1/20000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human
Host Species	Rabbit
Immunogen	Synthesized peptide derived from 2'-5'-oligoadenylate synthase 2 at AA range: 61-110
Specificity	OAS2 Polyclonal Antibody detects endogenous levels of OAS2 protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	OAS2
Accession No.	P29728 E9Q9A9 Q5MYU0
Alternate Names	OAS2; 2'-5'-oligoadenylate synthase 2; (2-5')oligo(A) synthase 2; 2-5A synthase 2; p69 OAS / p71 OAS; p69OAS / p71OAS

Antibody Specification

Description	2'-5'-oligoadenylate synthetase 2(OAS2) Homo sapiens This gene encodes a member of the 2-5A synthetase family, essential proteins involved in the innate immune response to viral infection. The encoded protein is induced by interferons and uses adenosine triphosphate in 2'-specific nucleotidyl transfer reactions to synthesize 2',5'-oligoadenylates (2-5As). These molecules activate latent RNase L, which results in viral RNA degradation and the inhibition of viral replication. The three known members of this gene family are located in a cluster on chromosome 12. Alternatively spliced transcript variants encoding different isoforms have been described. [provided by RefSeq, Jul 2008],
Protein Expression	Blood,Colon,Testis,Uterus,
Subcellular Localization	nucleus,cytoplasm,mitochondrion,endoplasmic reticulum,cytosol,membrane,intracellular membrane-bounded organelle,perinuclear region of cytoplasm,
Protein Function	catalytic activity: Binds double-stranded RNA and polymerizes ATP into PPP(A2'P5'A)N oligomers, which activate the latent RNase L that, when activated, cleaves single-stranded RNAs.,function: May play a role in mediating resistance to virus infection, control of cell growth, differentiation, and apoptosis.,induction: By interferons.,similarity: Belongs to the 2-5A synthetase family.,subcellular location: Associated with different subcellular fractions such as mitochondrial, nuclear, and rough/smooth microsomal fractions.,subunit: Homodimer.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.