

Immunotag™ Probetacellulin Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT5642
Product Description	Immunotag™ Probetacellulin Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	Probetacellulin
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human
Host Species	Rabbit
Immunogen	The antiserum was produced against synthesized peptide derived from the Internal region of human BTC. AA range:21-70
Specificity	Probetacellulin Polyclonal Antibody detects endogenous levels of Probetacellulin protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	BTC
Accession No.	P35070 Q05928
Alternate Names	BTC; Probetacellulin

Antibody Specification

Description	betacellulin(BTC) Homo sapiens This gene encodes a member of the epidermal growth factor (EGF) family of proteins. Alternative splicing results in multiple transcript variants, at least one of which encodes a preproprotein that is proteolytically processed to generate the secreted growth factor. A secreted form and a membrane-anchored form of this protein bind to multiple different EGF receptors. This protein promotes pancreatic cell proliferation and insulin secretion, as well as retinal vascular permeability. Mutations in this gene may be associated with type 2 diabetes in human patients. [provided by RefSeq, Nov 2015],
Cell Pathway/ Category	ErbB_HER,
Protein Expression	Mammary gland,Ovary,
Subcellular Localization	extracellular region,extracellular space,intracellular,plasma membrane,integral component of membrane,
Protein Function	disease:Betacellulin from beta cells could play a role in the vascular complications associated with diabetes.,function:Potent mitogen for retinal pigment epithelial cells and vascular smooth muscle cells. The effects of betacellulin are probably mediated by the EGF receptor and other related receptors.,similarity:Contains 1 EGF-like domain.,subunit:Monomer.,tissue specificity:Synthesized in several tissues and tumor cells. Predominantly expressed in pancreas and small intestine.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.