

Immunotag™ Vitronectin Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT5706
Product Description	Immunotag™ Vitronectin Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	Vitronectin
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Reactive Species	Human
Host Species	Rabbit
Immunogen	The antiserum was produced against synthesized peptide derived from the Internal region of human VTN. AA range:50-100
Specificity	Vitronectin Polyclonal Antibody detects endogenous levels of Vitronectin protein.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	VTN
Accession No.	P04004 P29788
Alternate Names	VTN; Vitronectin; VN; S-protein; Serum-spreading factor; V75

Antibody Specification

Description	vitronectin(VTN) Homo sapiens The protein encoded by this gene is a member of the pexin family. It is found in serum and tissues and promotes cell adhesion and spreading, inhibits the membrane-damaging effect of the terminal cytolytic complement pathway, and binds to several serpin serine protease inhibitors. It is a secreted protein and exists in either a single chain form or a clipped, two chain form held together by a disulfide bond. [provided by RefSeq, Jul 2008],
Cell Pathway/ Category	Focal adhesion,ECM-receptor interaction,
Protein Expression	Epithelium,Liver,Lung,Mammary gland,Pituitary,Plasma,Platelet,
Subcellular Localization	extracellular region,basement membrane,extracellular space,Golgi lumen,extracellular matrix,rough endoplasmic reticulum lumen,extracellular exosome,alpha-v-beta3 integrin-vitronectin complex,blood microparticle,
Protein Function	domain:The SMB domain mediates interaction with SERPINE1/PAI1. The heparin-binding domain mediates interaction with insulin.,function:Somatomedin-B is a growth hormone-dependent serum factor with protease-inhibiting activity.,function:Vitronectin is a cell adhesion and spreading factor found in serum and tissues. Vitronectin interact with glycosaminoglycans and proteoglycans. Is recognized by certain members of the integrin family and serves as a cell-to-substrate adhesion molecule. Inhibitor of the membrane-damaging effect of the terminal cytolytic complement pathway.,PTM:It has been suggested that the active SMB domain may be permitted considerable disulfide bond heterogeneity or variability, thus two alternate disulfide patterns based on 3D structures are described with 1 disulfide bond conserved in both.,PTM:N- and O-glycosylated.,PTM:Phosphorylation on Thr-69 and Thr-76 favors cell adhesion and spreading.,PTM:Sulfated on 2 tyrosine residues.,similarity:Contains 1 SMB (somatomedin-B) domain.,similarity:Contains 4 hemopexin-like domains.,subunit:Exists in two forms: a single chain 75 kDa form (V75) and a clipped form composed of two chains (65 kDa and 10 kDa) (V65+V10) which are held together by a disulfide bond. Interacts with SERPINE1/PAI1 and insulin.,tissue specificity:Plasma.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.