

Immunotag™ TIMP-4 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT5799
Product Description	Immunotag™ TIMP-4 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	TIMP-4
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	WB 1:500-2000, ELISA 1:10000-20000
Concentration	1 mg/ml
Reactive Species	Human
Host Species	Rabbit
Immunogen	Synthesized peptide derived from TIMP-4 at AA range: 174-223
Specificity	TIMP-4 Polyclonal Antibody detects endogenous levels of TIMP-4
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	TIMP4
Accession No.	Q99727 Q9JHB3
Alternate Names	Metalloproteinase inhibitor 4 (Tissue inhibitor of metalloproteinases 4) (TIMP-4)
Description	TIMP metalloproteinase inhibitor 4(TIMP4) Homo sapiens This gene belongs to the TIMP gene family. The proteins encoded by this gene family are inhibitors of the matrix metalloproteinases, a group of peptidases involved in degradation of the extracellular matrix. The secreted, netrin domain-containing protein encoded by this gene is involved in regulation of platelet aggregation and recruitment and may play role in hormonal regulation and endometrial tissue remodeling. [provided by RefSeq, Jul 2008],

Antibody Specification

Protein Expression	Brain,Heart,
Subcellular Localization	proteinaceous extracellular matrix,extracellular space,sarcomere,
Protein Function	function:Complexes with metalloproteinases (such as collagenases) and irreversibly inactivates them. Known to act on MMP-1, MMP-2, MMP-3, MMP-7 and MMP-9.,similarity:Belongs to the protease inhibitor I35 (TIMP) family.,similarity:Contains 1 NTR domain.,tissue specificity:Abundant in heart and present at low levels in many other tissues.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.