

# Immunotag™ PENK Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT5829
Product Description	Immunotag™ PENK Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	PENK
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	WB 1:500-2000, ELISA 1:10000-20000
Concentration	1 mg/ml
Reactive Species	Human
Host Species	Rabbit
Immunogen	Synthetic peptide from human protein at AA range: 51-100
Specificity	The antibody detects endogenous PENK protein
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	PENK
Accession No.	P01210 P22005
Alternate Names	PENK

## Antibody Specification

Description	proenkephalin(PENK) Homo sapiens This gene encodes a preproprotein that is proteolytically processed to generate multiple protein products. These products include the pentapeptide opioids Met-enkephalin and Leu-enkephalin, which are stored in synaptic vesicles, then released into the synapse where they bind to mu- and delta-opioid receptors to modulate the perception of pain. Other non-opioid cleavage products may function in distinct biological activities. [provided by RefSeq, Jul 2015],
Protein Expression	Brain,Cerebellum,Testis,
Subcellular Localization	extracellular region,plasma membrane,dendrite,symmetric synapse,neuronal cell body,perikaryon,axon terminus,cell body fiber,
Protein Function	function:Met- and Leu-enkephalins compete with and mimic the effects of opiate drugs. They play a role in a number of physiologic functions, including pain perception and responses to stress. PENK(114-133) and PENK(237-258) increase glutamate release in the striatum. PENK(114-133) decreases GABA concentration in the striatum.,PTM:The N-terminal domain contains 6 conserved cysteines thought to be involved in disulfide bonding and/or processing.,similarity:Belongs to the opioid neuropeptide precursor family.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.