

## Immunotag™ NFAT5 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT5853
Product Description	Immunotag™ NFAT5 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	NFAT5
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,IHC-p,ELISA
Recommended Dilution	WB 1:500-2000,IHC-p 1:500-200, ELISA 1:10000-20000
Concentration	1 mg/ml
Reactive Species	Human,Rat,Mouse
Host Species	Rabbit
Immunogen	Synthetic peptide from human protein at AA range: 212-284
Specificity	The antibody detects endogenous NFAT5
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	NFAT5 KIAA0827 TONEBP
Accession No.	O94916 Q9WV30
Alternate Names	Nuclear factor of activated T-cells 5 (NF-AT5) (T-cell transcription factor NFAT5) (Tonicity-responsive enhancer-binding protein) (TonE-binding protein) (TonEBP)

## Antibody Specification

Description	nuclear factor of activated T-cells 5(NFAT5) Homo sapiens The product of this gene is a member of the nuclear factors of activated T cells family of transcription factors. Proteins belonging to this family play a central role in inducible gene transcription during the immune response. This protein regulates gene expression induced by osmotic stress in mammalian cells. Unlike monomeric members of this protein family, this protein exists as a homodimer and forms stable dimers with DNA elements. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],
Cell Pathway/ Category	WNT,WNT-T CELLAxon guidance,VEGF,Natural killer cell mediated cytotoxicity,T_Cell_Receptor,B_Cell_Antigen,
Protein Expression	Brain,PCR rescued clones,
Subcellular Localization	nucleus,nucleoplasm,cytoplasm,integral component of membrane,
Protein Function	Experimental confirmation may be lacking for some isoforms,function:Plays a role in the inducible expression of genes. Regulates hypertonicity-induced cellular accumulation of osmolytes.,similarity:Contains 1 RHD (Rel-like) domain.,subunit:Does not bind with Fos and Jun transcription factors. But might be capable of forming stable dimers with DNA elements.,tissue specificity:Highest levels in skeletal muscle, brain, heart and peripheral blood leukocytes. Also expressed in placenta, lung, liver, kidney, pancreas, spleen, thymus, prostate, testis, ovary, small intestine and colon.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.