

# Immunotag™ CD275 Polyclonal Antibody

Antibody Specification	
Catalog No.	ITT6038
Product Description	Immunotag™ CD275 Polyclonal Antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	CD275
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	IHC-p,ELISA
Recommended Dilution	IHC-p 1:50-200, ELISA 1:10000-20000
Concentration	1 mg/ml
Reactive Species	Human
Host Species	Rabbit
Immunogen	Synthetic peptide from human protein at AA range: 201-250
Specificity	The antibody detects endogenous CD275
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	ICOSLG B7H2 B7RP1 ICOSL KIAA0653
Accession No.	O75144 Q9JHJ8
Alternate Names	ICOS ligand (B7 homolog 2) (B7-H2) (B7-like protein GI50) (B7-related protein 1) (B7RP-1) (CD antigen CD275)

## Antibody Specification

Description	<p>alternative products:Additional isoforms seem to exist,function:Ligand for the T-cell-specific cell surface receptor ICOS. Acts as a costimulatory signal for T-cell proliferation and cytokine secretion; induces also B-cell proliferation and differentiation into plasma cells. Could play an important role in mediating local tissue responses to inflammatory conditions, as well as in modulating the secondary immune response by co-stimulating memory T-cell function.,induction:Constitutive expression is further enhanced by treatment with TNF-alpha in peripheral blood B-cells and monocytes, while it is decreased in dendritic cells.,sequence caution:The sequence differs from that shown in position 300 onward for unknown reason.,sequence caution:Translation N-terminally shortened.,similarity:Belongs to the immunoglobulin superfamily. BTN/MOG family.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 1 Ig-like V-type (immunoglobulin-like) domain.,tissue specificity:Isoform 1 is widely expressed (brain, heart, kidney, liver, lung, pancreas, placenta, skeletal muscle, bone marrow, colon, ovary, prostate, testis, lymph nodes, leukocytes, spleen, thymus and tonsil), while isoform 2 is detected only in lymph nodes, leukocytes and spleen. Expressed on activated monocytes and dendritic cells.,</p>
Cell Pathway/ Category	Cell adhesion molecules (CAMs),Intestinal immune network for IgA production,
Protein Expression	Adrenal gland,Brain,Colon,Dendritic cell,Leukocyte,Peripheral blood
Subcellular Localization	plasma membrane,integral component of membrane,extracellular exosome,
Protein Function	<p>Additional isoforms seem to exist,function:Ligand for the T-cell-specific cell surface receptor ICOS. Acts as a costimulatory signal for T-cell proliferation and cytokine secretion; induces also B-cell proliferation and differentiation into plasma cells. Could play an important role in mediating local tissue responses to inflammatory conditions, as well as in modulating the secondary immune response by co-stimulating memory T-cell function.,induction:Constitutive expression is further enhanced by treatment with TNF-alpha in peripheral blood B-cells and monocytes, while it is decreased in dendritic cells.,sequence caution:The sequence differs from that shown in position 300 onward for unknown reason.,sequence caution:Translation N-terminally shortened.,similarity:Belongs to the immunoglobulin superfamily. BTN/MOG family.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 1 Ig-like V-type (immunoglobulin-like) domain.,tissue specificity:Isoform 1 is widely expressed (brain, heart, kidney, liver, lung, pancreas, placenta, skeletal muscle, bone marrow, colon, ovary, prostate, testis, lymph nodes, leukocytes, spleen, thymus and tonsil), while isoform 2 is detected only in lymph nodes, leukocytes and spleen. Expressed on activated monocytes and dendritic cells.,</p>
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.