

Immunotag™ mTOR antibody

Antibody Specification	
Catalog No.	ITT6120
Product Description	Immunotag™ mTOR antibody
Size	50 µg, 100 µg
Conjugation	HRP, Biotin, FITC, Alexa Fluor® 350, Alexa Fluor® 405, Alexa Fluor® 488, Alexa Fluor® 555, Alexa Fluor® 594, Alexa Fluor® 647
IMPORTANT NOTE	This product is custom manufactured with a lead time of 3-4 weeks. Once in production, this item cannot be cancelled from an order and is not eligible for return.
Target Protein	mTOR
Clonality	Polyclonal
Storage/Stability	-20°C/1 year
Application	WB,ELISA
Recommended Dilution	WB 1:500-2000, ELISA 1:10000-20000
Concentration	1 mg/ml
Reactive Species	Human,Mouse,Rat
Host Species	Rabbit
Immunogen	Synthesized peptide derived from human mTOR
Specificity	This antibody detects endogenous levels of mTOR antibody
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Form	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Gene Name	MTOR FRAP FRAP1 FRAP2 RAFT1 RAPT1
Accession No.	P42345 Q9JLN9 P42346
Alternate Names	Serine/threonine-protein kinase mTOR (EC 2.7.11.1) (FK506-binding protein 12-rapamycin complex-associated protein 1) (FKBP12-rapamycin complex-associated protein) (Mammalian target of rapamycin) (mTOR) (Mechanistic target of rapamycin) (Rapamycin and FKBP12 target 1) (Rapamycin target protein 1)

Antibody Specification

Description	mechanistic target of rapamycin(MTOR) Homo sapiens The protein encoded by this gene belongs to a family of phosphatidylinositol kinase-related kinases. These kinases mediate cellular responses to stresses such as DNA damage and nutrient deprivation. This protein acts as the target for the cell-cycle arrest and immunosuppressive effects of the FKBP12-rapamycin complex. The ANGPTL7 gene is located in an intron of this gene. [provided by RefSeq, Sep 2008],
Cell Pathway/ Category	ErbB_HER,mTOR,Insulin_Receptor,Adipocytokine,Type II diabetes mellitus,Pathways in cancer,Glioma,Prostate cancer,Acute myeloid leukemia,
Protein Expression	B-cell,Brain,Cerebellum,
Subcellular Localization	Golgi membrane,nucleus,nucleoplasm,cytoplasm,mitochondrial outer membrane,lysosome,lysosomal membrane,endoplasmic reticulum membrane,cytosol,phosphatidylinositol 3-kinase complex,endomembrane system,membrane,P
Protein Function	function:Acts as the target for the cell-cycle arrest and immunosuppressive effects of the FKBP12-rapamycin complex. Part of the TORC2 complex which plays a critical role in AKT1 Ser-473 phosphorylation, and may modulate the phosphorylation of PKCA and regulate actin cytoskeleton organization.,similarity:Belongs to the PI3/PI4-kinase family.,similarity:Contains 1 FAT domain.,similarity:Contains 1 FATC domain.,similarity:Contains 1 PI3K/PI4K domain.,similarity:Contains 7 HEAT repeats.,subunit:Interacts with the FKBP12-rapamycin complex. Binds UBQLN1. Forms part of the mammalian target of rapamycin 2 complex (TORC2) comprised of FRAP1, GBL, PRR5, RICTOR and SIN. TORC2 does not bind to and is not sensitive to FKBP12-rapamycin. Binds directly to PRR5 and RICTOR within the TORC2 complex.,tissue specificity:Expressed in numerous tissues, with highest levels in testis.,
Usage	For Research Use Only! Not for diagnostic or therapeutic procedures.