

GenCRISPR™ SpCas9 Antibody (4A1), mAb, Mouse

Cat. No.: A01935-40**Host:** Mouse**Size:** 40 µg**Ig Subclass:** IgG2a,K**Clone:** 4A1**Immunogen:** Recombinant*Streptococcus pyogenes*

CRISPR/Cas9

UniProt Accession: Q99ZW2**Gene ID:** 901176**Purification:** Protein A affinity column**Conjugation:** Unconjugated**Version:** 10/16/2017

PRODUCT INFORMATION

Description

Clustered regularly interspaced short palindromic repeats (CRISPR)/CRISPR-associated (Cas) protein 9 system provides a robust and multiplexable genome editing tool, enabling researchers to precisely manipulate specific genomic elements, and facilitating the elucidation of target gene function in biology and diseases. CRISPR/Cas9 genome editing allows for double-stranded DNA breaks at specific sequences to efficiently disrupt, excise, mutate, insert, or replace genes. To date, the *Streptococcus pyogenes* Cas9 (SpCas9) has been used broadly to achieve efficient genome editing in a variety of species and cell types. It is important that the precision of transfection and the level of Cas9 expression should be controlled strictly during the editing processes by using specific anti-CRISPR/Cas9 antibodies.

Specificity

The product is specific for *Streptococcus pyogenes* CRISPR/Cas9. This antibody binds with recombinant *Streptococcus pyogenes* CRISPR/Cas9 protein in ELISA and Endogenous overexpressed *Streptococcus pyogenes* CRISPR/Cas9 in immunofluorescence, western blot.

Concentration

0.5 mg/ml, lyophilized with PBS, pH 7.4, containing 0.02% sodium azide. (It may be customized for bulk orders.)

Note

GenScript can customize this product per the customer's request including product size, buffer components, etc.

Reconstitution

Reconstitute the lyophilized antibody with deionized water (or equivalent) to a final concentration of 0.5 mg/ml.

Storage

The lyophilized product remains stable up to 1 year at -20 °C from date of receipt. Upon reconstitution, it can be stored for 2-3 weeks at 2-8 °C or for up to 12 months at -20 °C or below. Avoid repeated freezing and thawing cycles.

** For non-clinical research use only. **

860 Centennial Ave., Piscataway, NJ 08854, USA

Toll-Free: 1-877-436-7274 Tel: 1-732-885-9188 Fax: 1-732-210-0262 Email: product@genscript.com Web: www.genscript.com

Applications

Working concentrations for specific applications should be determined by the investigators. The appropriate concentration may be affected by secondary antibody affinity, antigen concentration, the sensitivity of the method of detection, temperature, the length of the incubations, and other factors. The suitability of this antibody for applications other than those listed below has not been determined. The following concentration ranges are recommended starting points for this product.


ELISA detection: 0.01-1 $\mu\text{g/ml}$

Immunofluorescence: 0.2-1 $\mu\text{g/ml}$

Western blot: 0.5-2 $\mu\text{g/ml}$

Other applications: user-optimized


Example



** For non-clinical research use only. **

860 Centennial Ave., Piscataway, NJ 08854, USA


Toll-Free: 1-877-436-7274 Tel: 1-732-885-9188 Fax: 1-732-210-0262 Email: product@genscript.com Web: www.genscript.com



** For non-clinical research use only. **

860 Centennial Ave., Piscataway, NJ 08854, USA


Toll-Free: 1-877-436-7274 Tel: 1-732-885-9188 Fax: 1-732-210-0262 Email: product@genscript.com Web: www.genscript.com



Standard curve of SpCas9 Sandwich ELISA. The SpCas9 Sandwich ELISA assay is developed by using **GenCRISPR™ SpCas9 Antibody (14B6)** (GenScript, A01936-40) and **GenCRISPR™ SpCas9 Antibody (4A1)** (GenScript, A01935-40) as capture and detection antibody, respectively. These two antibodies recognize different epitopes.

In this ELISA assay, **GenCRISPR™ SpCas9 Antibody (4A1)** (GenScript, A01935-40) was labeled with Biotin. GenScript can provide customized conjugation service for this product per customer's request.

The sensitivity is < 1 ng/ml and the detection range is 0-10 ng/ml.



Western Blot of recombinant *Streptococcus pyogenes* CRISPR/Cas9 protein with two independent antibodies: **GenCRISPR™ SpCas9 Antibody (4A1)** (GenScript, A01935-40) and **GenCRISPR™ SpCas9 Antibody (14B6)** (GenScript, A01936-40).

The correlated pattern indicates the high specificity of these two antibodies.

Lane 1: 50 ng SpCas9 recombinant protein

Lane 2: 25 ng SpCas9 recombinant protein

Lane 3: 10 ng SpCas9 recombinant protein

Lane 4: 5 ng SpCas9 recombinant protein

Lane 5: 50 ng SpCas9 recombinant protein

Lane 6: 25 ng SpCas9 recombinant protein

Lane 7: 10 ng SpCas9 recombinant protein

Lane 8: 5 ng SpCas9 recombinant protein

Primary Antibody:

Lane 1~4: **GenCRISPR™ SpCas9 Antibody (4A1)** (GenScript, A01935-40) 1 µg/ml


Lane 5~8: **GenCRISPR™ SpCas9 Antibody (14B6)** (GenScript, A01936-40) 1 µg/ml

Secondary Antibody:

Goat anti-Mouse IgG (H&L) [IRDye⁸⁰⁰] (Licor, 926-32211) 0.125 µg/ml

** For non-clinical research use only. **

860 Centennial Ave., Piscataway, NJ 08854, USA



Western Blot of HEK293 transfected with PX458 (pSpCas9(BB)-2A-GFP) or untransfected cell lysates with two independent antibodies: **GenCRISPR™ SpCas9 Antibody (4A1)** (GenScript, A01935-40) and **GenCRISPR™ SpCas9 Antibody (14B6)** (GenScript, A01936-40). The correlated pattern indicates the high specificity of these two antibodies.

Lane 1: 50 µg HEK293 transfected with PX458 cell Lysate

Lane 2: 50 µg Untransfected HEK293 cell Lysate

Lane 3: 40 ng SpCas9 recombinant protein

Lane 4: 50 µg HEK293 transfected with PX458 cell Lysate

Lane 5: 50 µg Untransfected HEK293 cell Lysate

Lane 6: 40 ng SpCas9 recombinant protein


Primary Antibody:

Lane 1~3: GenCRISPR™ SpCas9 Antibody (4A1) (GenScript, A01935-40) 1 µg/ml

Lane 4~6: GenCRISPR™ SpCas9 Antibody (14B6) (GenScript, A01936-40) 1 µg/ml

Secondary Antibody:

Goat anti-Mouse IgG (H&L) [IRDye⁸⁰⁰] (Licor,926-32211) 0.125 µg/ml



Western Blot of HEK293 transfected with PX458 (pSpCas9(BB)-2A-GFP) or untransfected cell lysates with **GenCRISPR™ SpCas9 Antibody (4A1)** (GenScript, A01935-40). The different concentration of cell lysates indicates the high affinity and sensitivity of the antibody.

Lane 1: 50 µg HEK293 transfected with PX458 cell Lysate

Lane 2: 25 µg HEK293 transfected with PX458 cell Lysate

Lane 3: 10 µg HEK293 transfected with PX458 cell Lysate

Lane 4: 50 µg Untransfected HEK293 cell Lysate

Primary Antibody:


GenCRISPR™ SpCas9 Antibody (4A1) (GenScript, A01935-40) 1 µg/ml

Secondary Antibody:

Goat anti-Mouse IgG (H&L) [IRDye⁸⁰⁰] (Licor,926-32211) 0.125 µg/ml

** For non-clinical research use only. **

860 Centennial Ave., Piscataway, NJ 08854, USA



Western Blot of HEK293 transfected with PX458 (pSpCas9(BB)-2A-GFP) or untransfected cell lysates with **GenCRISPR™ SpCas9 Antibody (4A1)** (GenScript, A01935-40). The different concentration of antibodies indicates the high affinity and sensitivity of the antibody.

Lane 1: 50 µg HEK293 transfected with PX458 cell Lysate

Lane 2: 50 µg Untransfected HEK293 cell Lysate

Lane 3: 40 ng SpCas9 recombinant protein

Lane 4: 50 µg HEK293 transfected with PX458 cell Lysate

Lane 5: 50 µg Untransfected HEK293 cell Lysate

Lane 6: 40 ng SpCas9 recombinant protein

Lane 7: 50 µg HEK293 transfected with PX458 cell Lysate

Lane 8: 50 µg Untransfected HEK293 cell Lysate

Lane 9: 40 ng SpCas9 recombinant protein

Primary Antibody:


Lane 1~3: GenCRISPR™ SpCas9 Antibody (4A1) (GenScript, A01935-40) 2 µg/ml

Lane 4~6: GenCRISPR™ SpCas9 Antibody (4A1) (GenScript, A01935-40) 1 µg/ml

Lane 7~9: GenCRISPR™ SpCas9 Antibody (4A1) (GenScript, A01935-40) 0.5 µg/ml

Secondary Antibody:

Goat anti-Mouse IgG (H&L) [IRDye⁸⁰⁰] (Licor, 926-32211) 0.125 µg/ml



Western Blot analysis of HEK293 transfected with various plasmids with **GenCRISPR™ SpCas9 Antibody (4A1)** (GenScript, A01935-40). The different HEK293 with transfected with various plasmids indicate the minimum cross reaction of the antibody.

Loading:

Lane 1: 50µg HEK293 cell lysate transfected with SaCas9(BB)-2A-GFP (J7RUA5, *Staphylococcus aureus*)

Lane 2: 50µg HEK293 cell lysate transfected with StCas9(BB)-2A-GFP (G3ECR1, *Streptococcus thermophiles*)

Lane 3: 50µg HEK293 cell lysate transfected with FnCpf1(BB)-2A-GFP (A0Q7Q2, *Francisella tularensis subsp. novicida (strain U112)*)

Lane 4: 50µg HEK293 cell lysate transfected with LbCpf1(BB)-2A-GFP (A0A182DWE3, *Lachnospiraceae bacterium ND2006*)

Lane 5: 50µg HEK293 cell lysate transfected with AsCpf1(BB)-2A-GFP (U2UMQ6, *Acidaminococcus sp. (strain BV3L6)*)

Lane 6: 50µg HEK293 cell lysate transfected with pSpCas9(BB)-2A-GFP (PX458,

** For non-clinical research use only. **

860 Centennial Ave., Piscataway, NJ 08854, USA

Q99ZW2, *Streptococcus pyogenes* serotype M1)
Lane 7: 50µg HEK293 cell lysate (Non-transfected)
Primary Antibody:
Lane 1~7: GenCRISPR™ SpCas9 Antibody (4A1) (GenScript, A01935-40) 1 µg/ml
Lane 1~7: THE™ Anti-α-tubulin mAb (mouse)(1F4) 0.5µg/ml
Secondary antibody:
Goat anti-Mouse IgG (H&L) [IRDye⁸⁰⁰] (Licor, 926-32211) 0.125 µg/ml

** For non-clinical research use only. **

860 Centennial Ave., Piscataway, NJ 08854, USA

Toll-Free: 1-877-436-7274 Tel: 1-732-885-9188 Fax: 1-732-210-0262 Email: product@genscript.com Web: www.genscript.com