

DATASHEET

Version 20181206

IL-1RA, Human

Cat. No.: Z02700-100

Size: 100.0 ug

Synonyms: Interleukin-1 receptor antagonist (IL-1ra), Human;

Description:

Interleukin-1 receptor antagonist (IL-1ra) is a member of the IL-1 family. Endogenous IL-1ra is produced in numerous animal disease models as well as in human autoimmune and chronic inflammatory diseases. It binds to IL-1 receptors in competition with IL-1, but does not elicit intracellular response from this binding. Its role in counteracting the proinflammatory effects of IL-1 is being studied by numerous research groups. IL-4 and IL-13 have been shown to amplify the stimulatory effect of IL-1-beta on the production of soluble and intracellular forms of IL-1ra. The regulated expression of IL-1ra in various cell types has been shown to be influenced by cytokines. In synovial fibroblasts the synthesis of IL-1ra is markedly enhanced by IL-1, TNF-alpha, or PDGF.

Amino Acid Sequence:

```
00001 RPSGRKSSKM QAFRIWDVNQ KTFYLRNNQL VAGYLQGPV
00041 NLEEKIDVVP IEPHALFLGI HGGKMCLSCV KSGDETRLQL
00081 EAVNITDLSE NRKQDKRFAF IRSDSGPTTS FESAACPGWF
00121 LCTAMEADQP VSLTNMPDEG VMVTKFYFQE DE
```

Source: *E. coli*

Species: Human

Biological Activity: Fully biologically active when compared to standard. The ED₅₀ as determined by inhibiting IL-1α-dependent proliferation of murine D10S cells is less than 40 ng/ml, corresponding to a specific activity of $> 2.5 \times 10^4$ IU/mg in the presence of 50 pg/ml rHuIL-1α.

Molecular Weight: Approximately 17.1 kDa, a single non-glycosylated polypeptide chain containing 152 amino acids.

Formulation: Lyophilized from a 0.2 μm filtered concentrated solution in PBS, pH 7.4.

Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Reconstitution: We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.

Purity: > 96 % by SDS-PAGE and HPLC analyses.

Endotoxin Level: Less than 1 EU/μg of rHuIL-1RA as determined by LAL method.

Storage: This lyophilized preparation is stable at 2-8 °C, but should be kept at -20 °C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8 °C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20 °C to -70 °C. Avoid repeated freeze/thaw cycles.