

DATASHEET Version 20181206

## **BAFF-R**, Human

Cat. No.: Z02725-50

**Size**: 50.0 ug

Synonyms: BAFF Receptor (BAFF-R), Human;

## **Description:**

BAFF Receptor (BAFF-R), a member of the TNFR superfamily, is highly expressed in spleen, lymph node, and resting B cells and to some extent in activated B cells, resting CD4+ cells and peripheral blood leukocytes. BAFF-R is a type III transmembrane protein that binds with high specificity to BAFF (TNFSF13B). BAFF-R/BAFF signaling plays a critical role in B cell survival and maturation.

## Amino Acid Sequence:

00001 MRRGPRSLRG RDAPAPTPCV PAECFDLLVR HCVACGLLRT 00041 PRPKPAGASS PAPRTALQPQ ESVGAGAGEA ALPLPG Source: E. coli

Species: Human

**Biological Activity**: Fully biologically active when compared to standard. The  $ED_{50}$  as determined by its ability to block BAFF induced mouse splenocyte survival is 1.0-5.0 µg/ml in the presence of 1.0 µg/ml of rHuBAFF.

**Molecular Weight**: Approximately 7.8 kDa, a single non-glycosylated polypeptide chain containing 76 amino acids.

Formulation: Lyophilized from a 0.2  $\mu m$  filtered concentrated solution in 20 mM PB, pH 8.0, 500 mM NaCl.

**Appearance**: Sterile Filtered White lyophilized (freeze-dried) powder.

**Reconstitution**: We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at  $\leq$  -20 °C. Further dilutions should be made in appropriate buffered solutions.

**Purity**: > 95 % by reduced SDS-PAGE analyses.

**Endotoxin Level**: Less than 1 EU/µg of rHuBAFF-R as determined by LAL method.

**Storage**: This lyophilized preparation is stable at 2-8 °C, but should be kept at -20 °C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8 °C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20 °C to -70 °C. Avoid repeated freeze/thaw cycles.

For Research Use Only