

## FGF-7, Mouse

**Cat. No.:** Z02777-10

**Size:** 10.0 ug

**Synonyms:** Fibroblast Growth Factor-7(FGF-7), Mouse ;

### Description:

Fibroblast Growth Factor-7 (FGF-7/KGF) is one of 23 known members of the FGF family. All FGFs have two conserved cysteine residues and share 30 - 50% sequence identity at the amino acid level. Proteins of this family play a central role during prenatal development and postnatal growth and regeneration of variety of tissues, by promoting cellular proliferation and differentiation. KGF-1/FGF-7 is a mitogen factor specific for epithelial cells and keratinocytes and signals through FGFR 2b. KGF-1/FGF-7 plays a role in kidney and lung development, angiogenesis, and wound healing.

### Amino Acid Sequence:

```
00001 CNDMSPEQTA TSVNCSSPER HTRSVDYMEG GDIVRRLFC
00041 RTQWYLRIDK RGKVKGTQEM KNSYNIMEIR TVAVGIVAIAK
00081 GVESEYYLAM NKEGKLYAKK ECNEDCNFKE LILENHNTY
00121 ASAKWTHSGG EMFVALNQKG IPVKGKTKK EQKTAHFLPM
00161 AIT
```

**Source:** *E. coli*

**Species:** Mouse

**Biological Activity:** Fully biologically active when compared to standard. The ED<sub>50</sub> as determined by thymidine uptake assay using FGF-receptors transfected BaF3 cells is less than 10 ng/ml, corresponding to a specific activity of > 1.0 × 10<sup>5</sup> IU/mg.

**Molecular Weight:** Approximately 18.7 kDa, a single, non-glycosylated polypeptide chain containing 163 amino acids.

**Formulation:** Lyophilized from a 0.2 µm filtered solution in 20 mM PB, pH 8.0, 1 M NaCl.

**Appearance:** Sterile Filtered White lyophilized (freeze-dried) powder.

**Reconstitution:** We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/ml. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.

**Purity:** > 96 % by SDS-PAGE and HPLC analyses.

**Endotoxin Level:** Less than 1 EU/µg of rMuKGF-1/FGF-7 as determined by LAL method.

**Storage:** This lyophilized preparation is stable at 2-8 °C, but should be kept at -20 °C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8 °C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20 °C to -70 °C. Avoid repeated freeze/thaw cycles.