

DATASHEET

Version 20181206

MIC-B, Human

Cat. No.: Z02801-50

Size: 50.0 ug

Synonyms: (NULL)

Description:

MIC-B (MHC class I chain-related gene B) is a trans-membrane glycoprotein that functions as a ligand for human NKG2D. A closely related protein, MIC-A, shares 85% amino acid identity with MIC-B. These 2 proteins are distantly related to the MHC class I proteins. MIC-A and MIC-B (MIC-A/B) possess three extracellular immunoglobulin-like domains, but have no capacity to bind peptide or interact with β 2-microglobulin. The genes encoding MIC-A/B are found within the major histocompatibility complex on human chromosome 6. The MIC-B locus is polymorphic with more than 15 recognized human alleles. MIC-A/B are minimally expressed on normal cells, but are frequently expressed on epithelial tumors and can be induced by bacterial and viral infections. MIC-A/B are ligands for NKG2D, an activating receptor expressed on NK cells, NKT cells, $\gamma\delta$ T cells, and CD8+ β 223 T cells. Recognition of MIC-A/B by NKG2D results in the activation of cytolytic activity and/or cytokine production by these effector cells. MIC-A/B recognition is involved in tumor surveillance, viral infections, and autoimmune diseases. The release of soluble forms of MIC-A/B from tumors down-regulates NKG2D surface expression on effector cells resulting in the impairment of anti-tumor immune response.

Amino Acid Sequence:

```
00001 AEPHSLRYNL MVLSDQESVQ SGFLAEGHLD GQPFRLYDRQ
00041 KRRAPQGGQW AEDVLGAKTW DTETEDLTEN GQDLRLTLTH
00081 IKDQKGGHLS LQEIRVCEIH EDSSTRGSRH FYYDGELFLS
00121 QNLETQESTV PQSSRAQTLA MNVTNFWKED AMTKTHYRA
00161 MQADCLQKLQ RYLKSGVAIR RTVPPMVNVT CSEVSEGNIT
00201 VTCRASSFYP RNITLTWRQD GVSLSHNTQQ WG
```

Source: *E. coli*

Species: Human

Biological Activity: Fully biologically active when compared to standard. The specific activity is determined by binding MICB antibody in ELISA.

Molecular Weight: Approximately 32.8 kDa, a single non-glycosylated polypeptide chain containing 287 amino acids.

Formulation: Lyophilized from a 0.2 μ m filtered concentrated solution in 20 mM Tris, 150 mM NaCl, pH 8.0.

Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Reconstitution: We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/ml. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.

Purity: > 95 % by SDS-PAGE and HPLC analyses.

Endotoxin Level: Less than 1 EU/ μ g of rHuMIC-B as determined by LAL method.

Storage: This lyophilized preparation is stable at 2-8 °C, but should be kept at -20 °C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8 °C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20 °C to -70 °C. Avoid repeated freeze/thaw cycles.