

DATASHEET

Version 20181206

MCP-4/CCL13, Human**Cat. No.:** Z02836-20**Size:** 20.0 ug**Synonyms:** MCP-4/CCL13, Human;**Description:**

CCL13 is a chemoattractant for monocytes and eosinophils, and activates basophils. In addition, it has been reported to be chemotactic for CD4+ and CD8+ T cells, with an activity almost equivalent to that of MCP-3. The bioactivities of CCL13 is most likely mediated by the CC chemokine receptors CCR-2 and CCR-3, both of which have been shown to bind CCL13.

Amino Acid Sequence:

00001 QPDALNVPST CCFTFSSKKI SLQRLKSYVI TTSRCPQKAV
00041 IFRTKLGKEI CADPKEKVVQ NYMKHLGRKA HTLKT

Source: *E. coli***Species:** Human

Biological Activity: Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human monocytes is in a concentration of 10-100 ng/ml.

Molecular Weight: Approximately 8.6 kDa, a single non-glycosylated polypeptide chain containing 75 amino acids.

Formulation: Lyophilized from a 0.2 µm filtered concentrated solution in 20 mM PB, pH 7.4, 130 mM NaCl.

Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Reconstitution: We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20°C. Further dilutions should be made in appropriate buffered solutions.

Purity: > 96 % by SDS-PAGE and HPLC analyses.

Endotoxin Level: Less than 1 EU/µg of rHuMCP-4/CCL13 as determined by LAL method.

Storage: This lyophilized preparation is stable at 2-8 °C, but should be kept at -20 °C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8 °C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20 °C to -70 °C. Avoid repeated freeze/thaw cycles.