

DATASHEET
Version 20181206**MDC/CCL22, Mouse****Cat. No.:** Z02856-20**Size:** 20.0 ug**Synonyms:** MDC/CCL22, Mouse;**Description:**

MDC is a CC chemokine that is produced in B cells, macrophages, monocyte-derived dendritic cells, activated NK cells and CD4 T cells. It signals through the CCR4 receptor. MDC chemoattracts monocytes, dendritic cells and NK cells and exerts HIV suppressive activity.

Amino Acid Sequence:

00001 GPYGANVEDS ICCQDYIRHP LPSRLVKEFF WTSKSCRKPG
00041 VVLITVKNRD ICADPRQVWV KLLHLKLS

Source: *E. coli***Species:** Mouse

Biological Activity: Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human activated lymphocytes is in a concentration range of 10-100 ng/ml.

Molecular Weight: Approximately 7.8 kDa, a single, non-glycosylated polypeptide chain containing 68 amino acids.

Formulation: Lyophilized from a 0.2 µm filtered concentrated solution in 20 mM PB, pH 7.4, 150 mM NaCl.

Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Reconstitution: We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.

Purity: > 97 % by SDS-PAGE and HPLC analyses.

Endotoxin Level: Less than 1 EU/µg of rMuMDC/CCL22 as determined by LAL method.

Storage: This lyophilized preparation is stable at 2-8 °C, but should be kept at -20 °C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8 °C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20 °C to -70 °C. Avoid repeated freeze/thaw cycles.