

DATASHEET

Version 20181206

M-CSF, Human**Cat. No.:** Z02914-10**Size:** 10.0 ug**Synonyms:** Macrophage Colony Stimulating Factor, CSF-1, Lanimostim, MCSF, MGC31930, M-CSF.**Description:**

Macrophage Colony-Stimulating Factor 1 (M-CSF), involved especially in monocytopoiesis,^[1] is a hematopoietic growth factor. In mammals, it exists three isoforms, which invariably share an N-terminal 32-aa signal peptide, a 149-residue growth factor domain, a 21-residue transmembrane region and a 37-aa cytoplasmic tail^[2]. The biological activity of human M-CSF is maintained within the 149-aa growth factor domain^[3], and it is only active in the disulfide-linked dimeric form^[4], which is bonded at Cys63.

Recombinant human Macrophage Colony-Stimulating Factor 1 (rhM-CSF) produced in *E. coli* is a disulfide-linked homodimer containing two non-glycosylated polypeptide chains of 159 amino acids each. A fully biologically active molecule, rhM-CSF has a molecular mass of 28 kDa analyzed by non-reducing SDS-PAGE and is obtained by proprietary refolding and chromatographic techniques at GenScript.

Amino Acid Sequence:

```
00001 MEEVSEYCSH MIGSGHLQSL QRLIDSQMET SCQITFEFVD
00041 QEQLKDPVCY LKKAFLLVQD IMEDTMRFRD NTPNAIAIVQ
00081 LQELSLRLKS CFTKDYEEHD KACVRTFYET PLQLLEKVKV
00121 VFNETKNLLD KDNIFSKNC NNSFAECSSQ GHERQSEGS
```

Source: *E. coli***Species:** Human**Biological Activity:** ED₅₀ of 1 - 3 ng/ml, measured by cell proliferation assay of M-NFS-60, corresponding to a specific activity of 3.3 x 10⁵ - 1 x 10⁶ units/mg.**Molecular Weight:** 28 kDa, observed by non-reducing SDS-PAGE.**Formulation:** Lyophilized after extensive dialysis against 50 mM Tris-HCl, pH 8.0.**Reconstitution:** Reconstituted in ddH₂O or PBS or Tris-HCl, pH 8.0 at 100 µg/ml.**Purity:** > 95% as analyzed by non-reducing SDS-PAGE.**Endotoxin Level:** <1 EU/µg, determined by LAL method.**Storage:** Lyophilized recombinant human Macrophage Colony-Stimulating Factor 1 (rhM-CSF) remains stable up to 6 months at lower than -70°C from date of receipt. Upon reconstitution, rhM-CSF should be stable up to 2 weeks at 4°C or up to 3 months at -20°C.