

**DATASHEET**  
Version 20181206**IL-4, Porcine****Cat. No.:** Z02928-50**Size:** 50.0 ug**Synonyms:** Interleukin-4, BCGF, BCDF, B cell stimulating factor, BSF-1, Lymphocyte stimulatory factor 1, IL-4, MGC79402, Binetrakin, Pittrakina**Description:**

Interleukin-4 (IL-4) is a pleiotropic cytokine regulates diverse T and B cell responses including cell proliferation, survival, and gene expression. It has important effects on the growth and differentiation of different immunologically competent cells. Interleukin-4 is produced by mast cells, T cells, and bone marrow stromal cells [1]. IL-4 regulates the differentiation of native CD4+ T cells (Th0 cells) into helper Th2 cells, and regulates the immunoglobulin class switching to the IgG1 and IgE isotypes. IL-4 has numerous important biological functions including stimulating B-cells activation, T-cell proliferation and CD4+ T-cells differentiation to Th2 cells. It is a key regulator in hormone control and adaptive immunity [2]. IL-4 also plays a major role in inflammation response and wound repair via activation of macrophage into M2 cells [3]. IL-4 is stabilized by three disulphide bonds forming a compact globular protein structure [4]. Four alpha-helix bundle with left-handed twist is dominated half of the protein structure with 2 overhand connections and fall into a 2-stranded anti-parallel beta sheet [5].

**Amino Acid Sequence:**

```
00001 HKCDITLQEI IKTLNLTAR KNSCMELPVT DVFAAPENTT
00041 EKETFCEAST VLRHIYRHHT CMKSLLSGLD RNLSSMANMT
00081 CSVHEAKKST LKDFLERLKT IMKEKYSKC
```

**Source:** CHO**Species:** Porcine**Biological Activity:** ED<sub>50</sub> < 0.25 ng/ml, measured in a cell proliferation assay using TF-1 human erythroleukemic cells, corresponding to a specific activity of >4 x 10<sup>6</sup> units/mg**Molecular Weight:** 15kDa, observed by reducing SDS-PAGE**Formulation:** Lyophilized after extensive dialysis against PBS.**Reconstitution:** Reconstituted in ddH<sub>2</sub>O or PBS at 100 µg/ml.**Purity:** > 95% as analyzed by SDS-PAGE and HPLC.**Endotoxin Level:** <0.2 EU/µg, determined by LAL method.**Storage:** Lyophilized recombinant porcine Interleukin 4 (IL-4) remains stable up to 6 months at lower than -70°C from date of receipt. Upon reconstitution, rIL-4 should be stable up to 1 week at 4°C or up to 2 months at -20°C.