

DATASHEET
Version 20181206**MDC/CCL22, Rat****Cat. No.:** Z02959-5**Size:** 5.0 ug

Synonyms: C-C motif chemokine 22, Small-inducible cytokine A22, Macrophage-derived chemokine, MDC(1-69), Stimulated T-cell chemotactic protein 1, CC chemokine STCP-1, CCL22, MDC, SCYA22, ABCD-1, DC/B-CK, MGC34554, A-152E5.1, CC chemokine ABCD-1, Activated B and dendritic cell-derived, DCBCK.

Description:

Macrophage-Derived Chemokine/CCL22 (MDC) is a CC chemokine that is produced in B cells, macrophages, monocyte-derived dendritic cells, activated NK cells and CD4 T cells. It signals through the CCR4 receptor. MDC chemoattracts monocytes, dendritic cells and NK cells and exerts HIV suppressive activity.

Amino Acid Sequence:

00001 GPYGANVEDS ICCQDYIRHP LPPRFVKEFY WTSKSCRKPG
00041 VVLITIKNRD ICADPRMLWV KKILHKLA

Source: *E. coli***Species:** Rat

Biological Activity: Fully biologically active when compared to standard. The biologically active determined by a chemotaxis bioassay using human T-lymphocytes is in a concentration range of 10-100 ng/ml.

Molecular Weight: Approximately 7.9 kDa, a single, non-glycosylated polypeptide chain containing 68 amino acids.

Formulation: Lyophilized from a 0.2 µm filtered concentrated solution in 2 × PBS, pH7.4.

Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Reconstitution: We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/ml. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.

Purity: > 96 % by SDS-PAGE and HPLC analyses.

Endotoxin Level: Less than 1 EU/µg of rRt-MDC/CCL22 as determined by LAL method.

Storage: This lyophilized preparation is stable at 2-8 °C, but should be kept at -20 °C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8 °C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20 °C to -70 °C. Avoid repeated freeze/thaw cycles.