

DATASHEET
Version 20181206**IL-1 β , Mouse****Cat. No.:** Z02985-50**Size:** 50.0 ug

Synonyms: Catabolin, Lymphocyte-activating factor (LAF), Endogenous Pyrogen (EP), Leukocyte Endogenous Mediator (LEM), Mononuclear Cell Factor (MCF)

Description:

Interleukin-1 Beta (IL-1 β) is a proinflammatory cytokine produced in a variety of cells including monocytes, tissue macrophages, keratinocytes and other epithelial cells. Both IL-1 alpha and IL-1 beta binds to the same receptor and has similar if not identical biological properties. These cytokines have a broad range of activities including, stimulation of thymocyte proliferation, by inducing IL-2 release, B-cell maturation and proliferation, mitogenic FGF-like activity and the ability to stimulate the release of prostaglandin and collagenase from synovial cells. However, whereas IL-1 beta is a secreted cytokine, IL-1 alpha is predominantly a cell-associated cytokine.

Recombinant mouse Interleukin-1 Beta(rmIL-1 β) produced in *E. coli* is a single non-glycosylated polypeptide chain containing 153 amino acids. A fully biologically active molecule, rmIL-1 β is obtained by proprietary chromatographic techniques at GenScript, with an apparent molecular weight of 17.5kDa analyzed by reducing SDS-PAGE.

Amino Acid Sequence:

```
00001 MVPIRQLHYR LRDEQQKSLV LSDPYELKAL HLNQQNINQQ
00041 VIFSMSFVQG EPSNDKIPVA LGLKGKNLYL SCVMKDGTP
00081 LQLESVDPKQ YPKKKMEKRF VFNKIEVKSK VEFESAEPN
00121 WYISTSQAEH KPVFLGNNSG QDIIDFTMES VSS
```

Source: *E. coli***Species:** Mouse

Biological Activity: ED₅₀<2.0 pg/ml, measured by the dose-dependent stimulation of mouse D10S cells, corresponding to a specific activity of > 5.0×10⁸ units/mg.

Molecular Weight: 17.5kDa, observed by reducing SDS-PAGE.

Formulation: Lyophilized after extensive dialysis against PBS.

Reconstitution: Reconstituted in ddH₂O or PBS at 100 µg/ml.

Purity: > 95% by SDS-PAGE and HPLC analyses.

Endotoxin Level: <0.2 EU/µg, determined by LAL method.

Storage: Lyophilized recombinant mouse Interleukin-1 Beta(rmIL-1 β) remains stable up to 6 months at lower than -70°C from date of receipt. Upon reconstitution, rmIL-1 β should be stable up to 2 weeks at 4°C or up to 3 months at -20°C.