

DATASHEET Version 20181206

IL-17A, Mouse

Cat. No.: Z03031-50

Size: 50.0 ug

Synonyms: IL-17A, CTLA-8

Description:

Interleukin-17A, (also known as CTLA-8) is a T cell-expressed pleiotropic cytokine that exhibits a high degree of homology to a protein encoded by the ORF13 gene of herpesvirus Saimiri. cDNA clones encoding IL-17 have been isolated from activated rat, mouse and human T cells. IL-17 represents a family of structurally-related cytokines that share a highly conserved C-terminal region but differ from one another in their N-terminal regions and in their distinct biological roles. The six known members of this family, IL-17A through IL-17F, are secreted as homodimers.

Recombinant murine IL-17A is a 15-20kDa glycosylated cytokine of 133 amino acid polypeptide chain that plays an important role in antimicrobial and chronic inflammation. IL-17 exhibits multiple biological activities on a variety of cells including the induction of IL-6 and IL-8 production in fibroblasts, the enhancement of surface expression of ICAM-1 in fibroblasts, activation of NF-kB and costimulation of T cell proliferation.

Amino Acid Sequence:

00001 AAIIPQSSAC PNTEAKDFLQ NVKVNLKVFN SLGAKVSSRR 00041 PSDYLNRSTS PWTLHRNEDP DRYPSVIWEA QCRHQRCVNA 00081 EGKLDHHMNS VLIQQEILVL KREPESCPFT FRVEKMLVGV 00121 GCTCVASIVR QAA Source: CHO Species: Mouse

Biological Activity: Measured by its ability to induce IL-1a, IL-4 and IL-6 production by primary mouse splenocytes.

Molecular Weight: 15-22 kDa, observed by reducing SDS-PAGE.

Formulation: Lyophilized after extensive dialysis against PBS.

Reconstitution: Reconstituted in ddH₂O or PBS at 100 µg/ml.

Purity: > 95% as analyzed by SDS-PAGE.

Endotoxin Level: < 0.2 EU/µg, determined by LAL method.

Storage: Lyophilized recombinant Murine Interleukin-17A (IL-17A) remains stable up to 6 months at lower than -70°C from date of receipt. Upon reconstitution, recombinant murine IL-17A should be stable up to 1 week at 4°C or up to 2 months at -20°C.