

**DATASHEET**  
Version 20181206**GRO/MGSA/CXCL1, Human****Cat. No.:** Z03079-25**Size:** 25.0 ug

**Synonyms:** Growth Regulated Protein/Melanoma Growth Stimulatory Activity, GRO  $\alpha$ , MGSA  $\alpha$ , CXCL1, NAP-3, GRO1, KC (murine) CINC (rat)

**Description:**

GRO/MGSA/CXCL1 has chemotactic activity for neutrophils. It may play a role in inflammation and exerts its effects on endothelial cells in an autocrine fashion. All three isoforms of GRO are CXC chemokines that can signal through the CXCR1 or CXCR2 receptors. GRO expression is inducible by serum or PDGF and/or by a variety of inflammatory mediators, such as IL-1 and TNF, in monocytes, fibroblasts, melanocytes and epithelial cells. In certain tumor cell lines, GRO is expressed constitutively.

**Amino Acid Sequence:**

00001 ASVATELRQ CLQTLQGIHP KNIQSVNVKS PGPCHAQTEV  
00041 IATLKNRKA CLNPASPIVK KIIEKMLNSD KSN

**Source:** CHO**Species:** Human**Biological Activity:** Active, measured in a functional assay using HUVEC cells.**Molecular Weight:** 7.8 kDa, observed by reducing SDS-PAGE.**Formulation:** Lyophilized after extensive dialysis against PBS.**Reconstitution:** Reconstituted in ddH<sub>2</sub>O or PBS at 100  $\mu$ g/ml.**Purity:** > 95% as analyzed by SDS-PAGE.**Endotoxin Level:** < 0.2 EU/ $\mu$ g, determined by LAL method.

**Storage:** Lyophilized recombinant Human GRO/MGSA/CXCL1 remains stable up to 6 months at lower than -70°C from date of receipt. Upon reconstitution, Human GRO/MGSA/CXCL1 should be stable up to 1 week at 4°C or up to 2 months at -20°C.