

DATASHEET
Version 20181206**OSM, Mouse****Cat. No.:** Z03143-50**Size:** 50.0 ug**Synonyms:** OSM**Description:**

Oncostatin M (OSM) is a multifunctional cytokine, and belongs to Interleukin-6 (IL-6) subfamily, which also includes IL-11, leukemia inhibitory factor (LIF), ciliary neurotropic factor, cardiotrophin-1, and novel neurotrophin-1. *In vivo*, OSM is secreted from activated T cells, monocytes, neutrophils, and endothelial cells. OSM is related to LIF, and shares a receptor with LIF in human. Human OSM can bind to gp130 and recruit OSM Receptor β or LIF Receptor β to form a ternary complex. OSM stimulates the growth of different types of cells, including megakaryocytes, fibroblasts, vascular endothelial cells, and T cells. OSM inhibits the proliferation of several cancer cell lines, such as solid tissue tumor cells, lung cancer cells, melanoma cells, and breast cancer cells. Recombinant Mouse Oncostatin M (rmOSM) produced in *E. coli* is a single non-glycosylated polypeptide chain containing 182 amino acids. A fully biologically active molecule, rmOSM has a molecular mass of 20.5 kDa analyzed by reducing SDS-PAGE and is obtained by proprietary chromatographic techniques at GenScript.

Amino Acid Sequence:

```
00001 MNRGCSNSSF QLLSQLQNA NLGTNTESLL EPYIRLQNLN
00041 TPDLRAACTQ HSAVFPSEDT LRQLSKPHFL STVYTTLDLV
00081 LYQLDALRQK FLKTPAFPKL DSARHNILGI RNNVFCMARL
00121 LNHSLEIPEP TQTDGASRS TTPDVFNFK IGSCGFLWGY
00161 HRFMGSGVRV FREWDDGSTR SR
```

Source: *E. coli***Species:** Mouse**Biological Activity:** ED₅₀ < 10 ng/mL, measured by a cell proliferation assay using Rat Embryo Brain cells, corresponding to a specific activity of > 1 × 10⁵ units/mg.**Molecular Weight:** 20.5 kDa, observed by reducing SDS-PAGE.**Formulation:** Lyophilized after extensive dialysis against PBS.**Reconstitution:** Reconstituted in ddH₂O or PBS at 100 µg/ml.**Purity:** > 95% as analyzed by SDS-PAGE and HPLC.**Endotoxin Level:** < 0.2 EU/µg, determined by LAL method.**Storage:** Lyophilized recombinant Mouse Oncostatin M (rmOSM) remains stable up to 6 months at lower than -70°C from date of receipt. Upon reconstitution, rmOSM remains stable up to 2 weeks at 4°C or up to 3 months at -20°C.