

IGF-II, Human

Cat. No.: Z03147-10

Size: 10.0 ug

Synonyms: Insulin-like Growth Factor-II, Somatomedin A

Description:

Insulin-like Growth Factor II (IGF-II) is a single chain 7 kDa polypeptide, and shares a high degree of homology with insulin. During circulation in vivo, IGF-II is complexed to high affinity binding proteins, IGF Binding Proteins (IGFBP), which act as circulating reservoirs, transport IGF-II, and prolong the half life of IGF-II. The receptors of IGF-II (IGFRs) are transmembrane tyrosine receptors, and are heterotetrameric consisting of two α -subunits and two β -subunits. IGFRs execute their role via intracellular signaling molecules, such as IRS, shc, and PI3K. The functions of IGF-II include promoting cell survival, growth, proliferation, differentiation and motility. In particular, IGF-II promotes proliferation, inhibits death, and stimulates transformation in breast cancer cells.

Recombinant human Insulin-like Growth Factor II (rhIGF-II) produced in *E. coli* is a single non-glycosylated polypeptide chain containing 68 amino acids. A fully biologically active molecule, rhIGF-II has a molecular mass of 7.6 kDa analyzed by reducing SDS-PAGE and is obtained by proprietary chromatographic techniques at GenScript.

Amino Acid Sequence:

00001 MAYRPSETLC GGELVDTLQF VCGDRGFYFS RPASRVSRRS
00041 RGIVECCFR SCDLALLETY CATPAKSE

Source: *E. coli*

Species: Human

Biological Activity: ED₅₀ < 20 ng/mL, measured by a cell proliferation assay using FDCP-1 cells, corresponding to a specific activity of > 5×10⁴ units/mg.

Molecular Weight: 7.6 kDa, observed by reducing SDS-PAGE.

Formulation: Lyophilized after extensive dialysis against 25mM Tris, pH8.0.

Reconstitution: Reconstituted in ddH₂O at 100 μ g/mL.

Purity: > 95% as analyzed by SDS-PAGE and HPLC.

Endotoxin Level: < 0.2 EU/ μ g, determined by LAL method.

Storage: Lyophilized recombinant human Insulin-like Growth Factor II (rhIGF-II) remains stable up to 6 months at lower than -70°C from date of receipt. Upon reconstitution, rhIGF-II remains stable up to 2 weeks at 4°C or up to 3 months at -20°C.