

BRD1 (561-668aa), His, Human

Cat. No.: Z03184-100

Size: 100.0 ug

Synonyms: BRL, Bromodomain containing 1, BRPF2

Description:

Bromodomain (BRD) is an extensive family of protein domains, originally identified in and named after the *Drosophila* protein Brahma. Members of BRD family share a conserved atypical left-handed four helix bundle structure, and specifically bind to ϵ -lysine acetylated proteins. It is well known that histone acetylation and methylation play a central role in epigenetics and are important for various gene transcription events, thus the acetyl-lysine binding property of BRDs make them suitable drug targets for epigenetics. Currently, there are 46 diverse human proteins containing 61 BRDs. These include histone acetyltransferases, ATP-dependent chromatin-remodeling complex proteins, and nuclear scaffold proteins. The main functions of BRDs *in vivo* include chromatin acetylation and deacetylation, nucleosome assembly and remodeling, and organizations of chromosome or chromatin domains.

Recombinant human BRD1 (561-668) with His tag produced in *E. coli* is a single, non-glycosylated polypeptide chain containing 115 amino acids. A fully biologically active molecule, BRD1 (561-668) has a molecular mass of 13.7 kDa analyzed by reducing SDS-PAGE and is obtained by proprietary chromatographic techniques at GenScript.

Amino Acid Sequence:

```
00001 MHHHHHHELRLTPLTVLLRSVLDQLQDKDPARIFAQPVSL
00041 KEVPDYLDHIKHPMDFATMRKRLEAQGYKNLHEFEEDFDL
00081 IIDNCMKYNA RDTVFYRAAVRLRDQGGVVL RQARR
```

Source: *E. coli*

Species: Human

Biological Activity: Assay condition: 50 mM HEPES, 50 mM NaCl, and 1 mM CHAPS, pH 7.5. Biotinylated acetyl-histone peptide substrate binding to Bromodomain reaction was performed by adding 200 nM substrate to serially diluted BRD, and incubation for 1 h at RT. Treatment with equivalent detection buffer (2 nM Eu-Anti-His and 40 nM U-light-SA in Assay buffer), and incubation for 1 h at RT subsequently followed.

Molecular Weight: 13.7 kDa, observed by reducing SDS-PAGE.

Formulation: Sterile liquid solution contains 25mM HEPES, pH7.5, 150mM NaCl, 5% glycerol, 0.5 mM TCEP. Frozen solution.

Purity: > 95% by SDS-PAGE and HPLC analysis.

Endotoxin Level: < 1EU/ μ g, determined by LAL method.

Storage: BRD1 (561-668) remains stable up to 6 months at -80°C from date of receipt. For maximum recovery of product, centrifuge original vial prior to removing the cap. Aliquot to avoid repeated thawing and freezing.