

DATASHEET
Version 20181206**IL-6, His, Mouse****Cat. No.:** Z03189-5**Size:** 5.0 ug**Synonyms:** 26 kDa protein, IFN- β 2, B cell differentiation factor (BCDF), BSF-2, HPGF, HSF, MGI-2**Description:**

Interleukin-6 (IL-6), also known as BSF-2, CDF and IFNB2, is a pleiotropic cytokine that participates in both pro-inflammatory and anti-inflammatory responses. It is produced mainly by T cells, macrophages, monocytes, endothelial cells and muscle cells. IL-6 binds to IL-6 receptor (IL-6R) to trigger the association of IL-6R with gp130, inducing signal transduction through JAKs and STATs. The biological functions of IL-6 are diverse. It stimulates B cell differentiation and antibody production, myeloma and plasmacytoma growth, and nerve cell differentiation. It also acts as a myokine, produced by muscle cells in response to muscle contraction and released into the blood stream to help break down fats and improve insulin resistance.

Amino Acid Sequence:

```
00001 FPTSQVRRGD FTEDTTPNRP VYTTSQVGGL ITHVLWEIVE
00041 MRKELCNGNS DCMNDDALA ENNLKLPEIQ RNDGCYQTGY
00081 NQEICLLKIS SGLLEYHSYL EYMKNNLKD NKKDKARVLQR
00121 DTETLIHIFN QEVKDLHKIV LPTPISNALL TDKLESQKEW
00161 LRTKTIQFIL KSLEEFVKVT LRSTRQTHHH HHH
```

Source: CHO**Species:** Mouse**Biological Activity:** ED₅₀ < 3 pg /ml, measured in a bioassay using 7TD1 cells.**Molecular Weight:** 24-26 kDa, observed by reducing SDS-PAGE.**Formulation:** Lyophilized after extensive dialysis against PBS.**Reconstitution:** Reconstituted in ddH₂O or PBS at 100 µg/ml.**Purity:** > 95% as analyzed by SDS-PAGE.**Endotoxin Level:** < 0.2 EU/µg, determined by LAL method.**Storage:** Lyophilized recombinant murine Interleukin-6 (IL-6), His, Murine remains stable up to 6 months at lower than -70°C from date of receipt. Upon reconstitution, murine Interleukin-6 (IL-6), His should be stable up to 1 week at 4°C or up to 2 months at -20°C.