

DATASHEET
Version 20181206**IL-3, His, Rat****Cat. No.:** Z03196-50**Size:** 50.0 ug**Synonyms:** MCGF (Mast cell growth factor), Multi-CSF, HCGF, P-cell stimulation factor**Description:**

Interleukin-3 (IL-3), also known as MCGF, Multi-CSF, HCGF and P-cell stimulation factor, belongs to the α -helix family of hematopoietic cytokines. It is produced by activated T-cells, mast cells and natural killer cells. IL-3 binds to the IL-3 receptor α subunit and recruits the signal-transducing common β chain. IL-3 induced signal transduction results in the survival, differentiation and proliferation of a variety of immune cells, such as macrophages, neutrophils, megakaryocytes, mast cells and hematopoietic stem cells. IL-3 often acts synergistically with other cytokines, such as IL-7, EPO, GM-CSF and IL-6, to exert its stimulative function.

Amino Acid Sequence:

00001 ISDRGSDAHH LLRTLDCRTI ALEILVKLPY PQVSGLNNSD
00041 DKANLRNSTL RRVNLDEFLK SQEEFDSQDT TDIKSKLQKL
00081 KCCIPAAASD SVLPGVYNKD LDDFKKKLRF YVIHLKDLQP
00121 VSVSRPPQPT SSSDNFRPMT VECHHHHHH

Source: CHO**Species:** Rat**Biological Activity:** ED₅₀ < 8 ng/ml, measured in a cell proliferation assay using NF S-60 cells.**Molecular Weight:** 22-34 kDa, observed by reducing SDS-PAGE.**Formulation:** Lyophilized after extensive dialysis against PBS.**Reconstitution:** Reconstituted in ddH₂O or PBS at 100 μ g/ml.**Purity:** > 95% as analyzed by SDS-PAGE.**Endotoxin Level:** < 0.2 EU/ μ g, determined by LAL method.**Storage:** Lyophilized recombinant rat Interleukin-3 (IL-3), His remains stable up to 6 months at lower than -70°C from date of receipt. Upon reconstitution, rat Interleukin-3 (IL-3), His should be stable up to 1 week at 4°C or up to 2 months at -20°C.