

## IL-7, His, Mouse

**Cat. No.:** Z03209-50

**Size:** 50.0 ug

**Synonyms:** Lymphopoietin 1(LP-1), pre-B cell factor

### Description:

Interleukin-7 (IL-7), also known as lymphopoietin 1 and pre-B cell factor, is a hematopoietic growth factor belonging to the IL-7/IL-9 family. It is produced by keratinocytes, dendritic cells, hepatocytes, neurons and epithelial cells. IL-7 binds and signals through IL-7 receptor, a heterodimer consisting of IL-7 receptor alpha and common gamma chain receptor. IL-7 plays a role in regulating early B cell and T cell development. It is also important for optimal dendritic cell-T cell interaction.

### Amino Acid Sequence:

```
00001 ECHIKDKEGK AYESVLMISI DELDKMTGTD SNCPNNEPNF
00041 FRKHVCDDTK EAAFLNRAAR KLKQFLKMNI SEEFNVHLLT
00081 VSQGTQTLVN CTSKEEKNVK EQKKNDACFL KRLLEIKTC
00121 WNKILKGSIH HHHHH
```

**Source:** CHO

**Species:** Mouse

**Biological Activity:** ED<sub>50</sub> < 0.4 ng/ml, measured in a cell proliferation assay using 2E8 cells.

**Molecular Weight:** 8-28 kDa, observed by reducing SDS-PAGE.

**Formulation:** Lyophilized after extensive dialysis against PBS.

**Reconstitution:** Reconstituted in ddH<sub>2</sub>O or PBS at 100 µg/ml.

**Purity:** > 95% as analyzed by SDS-PAGE.

**Endotoxin Level:** < 0.2 EU/µg, determined by LAL method.

**Storage:** Lyophilized recombinant murine Interleukin-7 (IL-7), His remains stable up to 6 months at lower than -70°C from date of receipt. Upon reconstitution, murine Interleukin-7 (IL-7), His should be stable up to 1 week at 4°C or up to 2 months at -20°C.