

**DATASHEET**  
Version 20181206**LD78-β/CCL3L1, Human****Cat. No.:** Z03226-25**Size:** 25.0 ug**Synonyms:** SCYA3L1**Description:**

LD78-beta/CCL3L1 is a proinflammatory chemokine and the isoform of Macrophage Inflammatory Protein-1 alpha (MIP-1 alpha). LD78-beta is secreted by most mature leukocytes, predominantly macrophages, and its major receptor is the G-protein coupled receptor CCR5, which is also the co-receptor used by the HIV-1 virus for cell entry. LD78-beta has superior antiviral activity and induces a variety of immune cells, particularly CD8<sup>+</sup> T cells and immature dendritic cells. LD78-beta attracts lymphocytes and macrophages to sites of inflammation and infection, and its functions are inhibited by Interleukin-4, Interleukin-10, and Interleukin-13. Importantly, the copy number variation of LD78-beta is associated with HIV susceptibility, indicating LD78-beta's critical role in the disease.

Recombinant human LD78-beta/CCL3L1 (rhLD78-beta) produced in *E.coli* is a single non-glycosylated polypeptide chain containing 70 amino acids. A fully biologically active molecule, rhLD78-beta has a molecular mass of 7.8 kDa analyzed by reducing SDS-PAGE and is obtained by chromatographic techniques at GenScript.

**Amino Acid Sequence:**

00001 APLAADTPTA CCFSYTSRQI PQNFIADYFE TSSQCSKPSV  
00041 IFLTKRGRQV CADPSEEWVQ KYVSDLELSA

**Source:** *E. coli***Species:** Human

**Biological Activity:** ED<sub>50</sub> < 0.4 µg/mL, measured by the FLIPR assay using CHO cells transfected with human CCR5, the receptor of human CCL3L1, corresponding to a specific activity of > 2.5×10<sup>3</sup> units/mg.

**Molecular Weight:** 7.8 kDa, observed by reducing SDS-PAGE.

**Formulation:** Lyophilized after extensive dialysis against PBS.

**Reconstitution:** Reconstituted in ddH<sub>2</sub>O at 100 µg/mL.

**Purity:** > 95% by SDS-PAGE and HPLC analysis.

**Endotoxin Level:** < 0.2 EU/µg, determined by LAL method.

**Storage:** Lyophilized recombinant human LD78-beta/CCL3L1 (rhLD78-beta) remains stable up to 6 months at lower than -70°C from date of receipt. Upon reconstitution, rhLD78-beta remains stable up to 2 weeks at 4°C or up to 3 months at -20°C.