

DATASHEET
Version 20181206**ENA-78/CXCL5 (5-78aa), Human****Cat. No.:** Z03261-25**Size:** 25.0 ug**Synonyms:** Epithelial Neutrophil Activating Peptide-78, CXCL5**Description:**

Epithelial cell derived neutrophil activating peptide (ENA78) also known as C-X-C motif chemokine 5 (CXCL5), is a small cytokine belonging to the CXC chemokine family. It is produced following stimulation of cells with the inflammatory cytokines interleukin-1 or tumor necrosis factor- α . Expression of CXCL5 has also been observed in eosinophils, and can be inhibited with the type II interferon, IFN- γ . This chemokine stimulates the chemotaxis of neutrophils possessing angiogenic properties. Full length CXCL5 (78 aa) is trimmed at the Nterminal end by cathepsin G and chymotrypsin to ENA-74 (74 aa) and ENA-70 (70aa), with the shortened forms showing increased potency relative to full length CXCL5. CXCL5 can signal through the CXCR2 receptor.

Recombinant human ENA-78/CXCL5 (5-78a.a.) produced in *E. coli* is a single non-glycosylated polypeptide chain containing 74 amino acids. A fully biologically active molecule, rh ENA-78/CXCL5 (5-78a.a.) has a molecular mass of 8 kDa analyzed by reducing SDS-PAGE and is obtained by chromatographic techniques at GenScript.

Amino Acid Sequence:

00001 AAVLRELRCV CLQTTQGVHP KMISNLQVFA IGPQCSKVEV
00041 VASLKNKGKEI CLDPEAPFLK KVIQKILDGG NKEN

Source: *E. coli***Species:** Human

Biological Activity: The EC₅₀ value of human ENA-78/CXCL5 (5-78a.a.) on Ca²⁺ mobilization assay in CHO-K1/ G α 15/hCXCR2 cells (human G α 15 and human CXCR2 stably expressed in CHO-K1 cells) is less than 50 ng/ml.

Molecular Weight: 8 kDa, observed by reducing SDS-PAGE.

Formulation: Lyophilized after extensive dialysis against PBS.

Reconstitution: Reconstituted in ddH₂O or PBS at 100 μ g/ml.

Purity: > 95% as analyzed by SDS-PAGE

Endotoxin Level: < 0.2 EU/ μ g, determined by LAL method.

Storage: Lyophilized recombinant human ENA-78/CXCL5 (5-78a.a.) remains stable up to 6 months at lower than -70°C from date of receipt. Upon reconstitution, human ENA-78/CXCL5 (5-78a.a.) should be stable up to 1 week at 4°C or up to 2 months at -20°C.