

DATASHEET Version 20181206

B7-H3 Fc Chimera, Human

Cat. No.: Z03426-100

Size: 100.0 ug

Synonyms: B7H3; B7-H3; B7H34Ig-B7-H3; B7-

H3B7; CD276ss

Description:

Human B7 homolog 3 (B7-H3), a member of the immunoglobulin superfamily, is also known CD276, which contains two Ig-like C2-type (immunoglobulin-like) domains and two Ig-like Vtype (immunoglobulin-like) domains. B7-H3 may participate in the regulation of T-cell-mediated immune response. B7-H3 also plays a protective role in tumor cells by inhibiting natural-killer mediated cell lysis as well as a role of marker for detection of neuroblastoma cells. Furthermore. B7-H3 is involved in the development of acute and chronic transplant rejection and in the regulation of lymphocytic activity at mucosal surfaces. Human B7-H3 does not bind any known members of the CD28 family of immunoreceptor. However, B7-H3 has been shown to bind an unidentified counter-receptor on activated T cells to co-stimulate the proliferation of CD4+ or CD8+ T cells. B7-H3 has also been found to enhance the induction of primary cytotoxic T lymphocytes and stimulate IFN-gamma production.

Recombinant Human B7-H3 produced in HEK293 cells is a polypeptide chain containing 450 amino acids with C-terminal human IgG1 Fc fragment. A fully biologically active molecule, rhB7-H3 has a molecular mass of 62-65 kDa analyzed by reducing SDS-PAGE and is obtained by chromatographic

techniques at GenScript.

Amino Acid Sequence:

00001 LEVQVPEDPV VALVGTDATL CCSFSPEPGF SLAQLNLIWQ
00041 LTDTKQLVHS FAEGQDQGSA YANRTALFPD LLAQGNASLR
00081 LQRVRVADEG SFTCFVSIRD FGSAAVSLQV AAPYSKPSMT
00121 LEPNKDLRPG DTVTITCSSY RGYPEAEVFW QDGQGVPLTG
00161 NVTTSQMANE QGLFDVHSVL RVVLGANGTY SCLVRNPVLQ
00201 QDAHGSVTIT GQPMTFP

Source: HEK293

Molecular Weight: 62-65 kDa, observed by reducing SDS-PAGE.

Formulation: Lyophilized from a $0.2~\mu m$ filtered solution in PBS.

Reconstitution: Reconstituted in ddH₂O or PBS at 100 µg/ml.

Purity: > 95% as analyzed by reducing SDS-PAGE.

Endotoxin Level: < 0.2 EU/µg, determined by LAL method.

Storage: Lyophilized recombinant B7-H3, Human remains stable up to 6 months at lower than -70°C from date of receipt. Upon reconstitution, Human B7-H3 should be stable up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.