

DATASHEET Version 20181206

TGF-β 1, Mouse

Cat. No.: Z03431-50

Size: 50.0 µg

Synonyms: TGF-beta-1; CED; DPD1; TGFB; TGF-b1; TGFB1; CEDLAP; latency-associated peptide; TGFbeta; TGF-beta 1 protein; transforming growth factor beta-1

Description:

Transforming growth factor beta 1 (TGFβ1) is the prototype of a growing superfamily of peptide growth factors and plays a prominent role in a variety of cellular processes, including cell-cycle progression, cell differentiation, reproductive function, development, motility, adhesion, neuronal growth, bone morphogenesis, wound healing, and immune surveillance. TGF-β1, TGF-β2 and TGF-β3 signal via the same heteromeric receptor complex, consisting of a ligand binding TGF-β receptor type II (TβR-II), and a TGF- β receptor type I (T β R-I). Signal transduction from the receptor to the nucleus is mediated via SMADs. TGF-β expression is found in cartilage, bone, teeth, muscle, heart, blood vessels, hematopoietic cells, lung, kidney, gut, liver, eye, ear, skin, and the nervous system.

Recombinant Mouse TGF- β 1 produced by a mammalian expression system is a polypeptide chain containing 112 amino acids. A fully biologically active molecule; rm TGF- β 1 a molecular mass of 12.8 kDa analyzed by reducing SDS-PAGE and is obtained by chromatographic techniques at GenScript.

Amino Acid Sequence:

Ala²⁷⁹-Ser³⁹⁰ (Accession #:P04202);

00001 ALDTNYCFSS TEKNCCVRQL YIDFRKDLGW KWIHEPKGYH 00041 ANFCLGPCPY IWSLDTQYSK VLALYNQHNP GASASPCCVP 00081 QALEPLPIVY YVGRKPKVEQ LSNMIVRSCK CS

Source: Human Cells

Biological Activity: ED50 <0.2 ng/ml, measured in a cell proliferation assay using mouse HT-2 cells.

Molecular Weight: 12.8 kDa; observed by reducing

SDS-PAGE.

Formulation: Lyophilized from a $0.2~\mu m$ filtered solution in of 4 mM HCl.

Reconstitution: Reconstituted in ddH_2O at 100 $\mu g/ml$.

Purity: > 95% as analyzed by SDS-PAGE.

Endotoxin Level: < 1 EU/ μ g; determined by LAL method.

Storage: Lyophilized recombinant TGF- β 1, Mouse remains stable up to 6 months at lower than -70°C from date of receipt. Upon reconstitution; Mouse TGF- β 1 should be stable up to 1 week at 4°C or up to 3 months at -20°C. For long term storage it is recommended that a carrier protein (example 0.1% BSA) be added. Avoid repeated freeze-thaw cycles.