

# Bulk Anti-Human CD19 (SJ25-C1) Antibody

Bulk anti-human CD19 antibody (SJ25-C1)

| <b>T</b> | •  | 4  | T  | Po 1    |
|----------|----|----|----|---------|
| PrΛ      | an | ∩t | KΔ | nefits: |
| 110      | uu | Lι | DU | ucnts.  |

ichorbio's bulk anti-human CD19 antibody (SJ25-C1) is manufactured in a cGMP compliant, ISO Quality Standard 9001:2015 facility. ichorbio's low endotoxin antibodies have half the endotoxin of comparable antibodies from <u>Bio X Cell</u> at less than 1.0 EU/mg. If ichorbio's low endotoxin antibodies are not low enough we also offer ultra low endotoxin antibodies which have even less endotoxin (0.5EU/mg) at an even higher purity (98% versus 95%). ichorbio: the best antibodies for *in vivo* research

| endotoxin antibodies which have even less endotoxin (0.5EU/mg) at an even higher purity (98% versus 95%). ichorbio: the best antibodies for <i>in vivo</i> research. |
|--|
| Target:  |
| CD19   |
| Clone:   |
| SJ25-C1  |
| Isotype:   |
| Mouse IgG1   |
| Other Names:   |
| B-lymphocyte antigen CD19, B-lymphocyte surface antigen B4, Differentiation antigen CD19, T-cell surface antigen Leu-12,   |
| Uniprot:   |
| <u>P15391</u>  |
| Host:  |
| Mouse  |
| Species Reactivity:  |
| Human  |
| Specificity:   |
| Bulk anti-human CD19 antibody (SJ25-C1) recognizes an epitope on Human CD19  |
| Purification Method:   |

This monoclonal antibody was purified using multi-step affinity chromatography methods such as Protein A or G depending on the species and isotype.



# **Antigen Distribution:**

Anti-Human CD19 recognizes a (Mr 90 kDa) B-cell surface glycoprotein which is present at all stages of maturation. Anti-CD19 does not react with T-cells, granulocytes, monocytes, erythrocytes or platelets. The CD19 antigen is expressed on approximately 5-25% of human peripheral blood lymphocytes and approximately 60% of splenic lymphocytes.

### Immunogen:

NALM1 + NALM16 cells

#### **Concentration:**

? 2.0 mg/ml

#### **Formulation:**

0.01 M phosphate buffered saline (PBS) pH 7.2, 150 mM NaCl with no carrier protein, potassium or preservatives added. BSA and Azide free.

## **Purity:**

>95% by SDS-PAGE and HPLC

>98% by SDS-PAGE and HPLC

#### **Endotoxin:**

0.5 EU/mg as determined by the LAL method

? 0.75 EU/mg as determined by the LAL method

### **Aggregation:**

Aggregation level ? 5%

Aggregation level ? 1%

#### **Storage:**

This antibody is stable for at least 4 weeks when stored at 2-8°C. For long term storage, aliquot in working volumes without diluting and store at -20°C or -80°C. Avoid repeated freeze thaw cycles.

# **Applications:**

Flow Cytometry

#### **Application Notes:**

Each investigator should determine their own optimal working dilution for specific applications.

#### Use:

ichor.bio // hello@ichor.bio

Products are for research use only. Not for use in diagnostic or the rapeutic procedures. ichorbio, ichorbio logo and all other trademarks are the property of ichorbio LTD © ichorbio 2023 Page 2



Products are for research use only.

**Isotype Control:** 

Mouse IgG1 Isotype Control for In Vivo - Low Endotoxin [HKSP] (ICH2247)

Antibodies against the same target:

Anti-CD19 In Vivo Antibody - Low Endotoxin [1D3] (ICH1052), Anti-CD19 In Vivo Antibody - Ultra Low Endotoxin [1D3] (ICH1052UL)