

Biotinylated Cynomolgus ACE2/ACEH Protein

Cat. No. ACE-CM401B

Description

Source	Recombinant Biotinylated Cynomolgus ACE2/ACEH Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus.
	It contains Gln18-Val739.
Accession	A0A2K5X283
Molecular Weight	The protein has a predicted MW of 86.5 kDa. Due to glycosylation, the protein migrates to 110-140 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC

Formulation and Storage

Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

ACE2 (Angiotensin I Converting Enzyme 2) is a Protein Coding gene. Diseases associated with ACE2 include Severe Acute Respiratory Syndrome and Neurogenic Hypertension. The protein encoded by this gene belongs to the angiotensin-converting enzyme family of dipeptidyl carboxydipeptidases and has considerable homology to human angiotensin 1 converting enzyme. This secreted protein catalyzes the cleavage of angiotensin I into angiotensin 1-9, and angiotensin II into the vasodilator angiotensin 1-7.

Assay Data

Bis-Tris PAGE



Biotinylated Cynomolgus ACE2 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.


SEC-HPLC

Biotinylated Cynomolgus ACE2/ACEH Protein

Cat. No. ACE-CM401B



Assay Data




The purity of Biotinylated Cynomolgus ACE2 is greater than 95% as determined by SEC-HPLC.

ELISA Data

Biotinylated Cynomolgus ACE2, His Tag ELISA

0.1µg SARS-COV-2 Spike RBD, hFc Tag Per Well



Immobilized SARS-COV-2 Spike RBD, hFc Tag at 1µg/ml (100µl/well) on the plate. Dose response curve for Biotinylated Cynomolgus ACE2, His Tag with the EC50 of 0.28µg/ml determined by ELISA.