# Biotinylated Human ANGPT2/Angiopoietin-2 Protein






| Description             |   |
|-------------------------|---|
| Source                  | Recombinant Biotinylated Human ANGPT2/Angiopoietin-2 Protein is expressed from HEK293 with His tag and Avi tag at the N-Terminus.   |
|                         | It contains Lys275-Phe496.  |
| Accession               | O15123-1  |
| Molecular<br>Weight     | The protein has a predicted MW of 28.4 kDa. Due to glycosylation, the protein migrates to 32-37 kDa based on Bis-Tris PAGE result.  |
| Endotoxin               | Less than 1 EU per μg by the LAL method.  |
| Purity                  | > 95% as determined by Bis-Tris PAGE  |
| Formulation and Storage |   |
| Formulation             | Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.  |
| Reconstitution          | Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.   |
| Storage                 | -20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.   |
| Background              |   |
|                         | Angiopoietin-2 (Ang-2; also ANGPT2) is a secreted glycoprotein that plays a complex role in angiogenesis and inflammation. Mature Ang-2 is 478 amino acids in length.Ang2 is widely expressed during development, but it is restricted postnatally to highly angiogenic tissues such as the placenta, ovaries, and uterus. It is particularly abundant in vascular endothelial cells (EC) where it is stored in intracellular Weibel Palade bodies. |

# **Assay Data**

## **Bis-Tris PAGE**



Biotinylated Human ANGPT2 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.


**ELISA Data** 

## **Assay Data**



## **Biotinylated Human ANGPT2, His Tag ELISA**


0.1µg Biotinylated Human ANGPT2, His Tag Per Well



Log Anti-ANGPT2 Antibody, hFc Tag Conc.(μg/ml)

Immobilized Biotinylated Human ANGPT2, His Tag at  $1\mu g/ml$  ( $100\mu l/well$ ) on the streptavidin precoated plate ( $5\mu g/ml$ ). Dose response curve for Anti-ANGPT2 Antibody, hFc Tag with the EC50 of 3.8ng/ml determined by ELISA (QC Test).

### **SPR Data**



Human TIE2, His Tag immobilized on CM5 Chip can bind Biotinylated Human ANGPT2, His Tag with an affinity constant of 2.93  $\mu$ M as determined in SPR assay (Biacore T200).