Human B2M/beta 2-Microglobulin Protein

ICI-resistant tumors.


Cat. No. B2M-HM101



Description	
Source	Recombinant Human B2M/beta 2-Microglobulin Protein is expressed from HEK293 with His tag at the C-Terminus.
	It contains Ile21-Met119.
Accession	P61769-1
Molecular Weight	The protein has a predicted MW of 12.8 kDa. Due to glycosylation, the protein migrates to 13-15 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC
Formulation and Storage	
Formulation	Lyophilized from 0.22μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Background	
	The genetic and functional analysis of β2-microglobulin (B2M), a component of the HLA class-I complex.Acquired homozygous loss of B2M that caused lack of cell-surface HLA Class I expression in the tumor and a matched patient-derived xenograft (PDX). Downregulation of B2M was also found in two additional PDXs established from

Assay Data

Bis-Tris PAGE




Human B2M on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Cat. No. B2M-HM101



Assay Data



The purity of Human B2M is greater than 95% as determined by SEC-HPLC.