

Biotinylated Human B7-2/CD86 Protein



Cat. No. B72-HM486B


Description	
Source	Recombinant Biotinylated Human B7-2/CD86 Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus. It contains Leu26-Pro247.
Accession	P42081-1
Molecular Weight	The protein has a predicted MW of 28.2 kDa. Due to glycosylation, the protein migrates to 55-70 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE

Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt.-80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background	
B7-1 and B7-2 are homologous costimulatory ligands expressed on the surface of antigen presenting cells (APCs). Binding of these molecules to the T cell costimulatory receptors, CD28 and CTLA-4, is essential for the activation and regulation of T cell immunity. B7-1 and B7-2 do not form hetero-oligomers, underscoring the biological relevance of dimeric and monomeric state of B7-1 and B7-2, respectively.	

Assay Data


Bis-Tris PAGE



ELISA Data

Biotinylated Human B7-2, His Tag ELISA

0.1µg Biotinylated Human B7-2, His Tag Per Well



Immobilized Biotinylated Human B7-2, His Tag at 1µg/ml (100µl/well) on the streptavidin precoated plate (5µg/ml). Dose response curve for Human CTLA-4, hFc Tag with the EC50 of 1.9ng/ml determined by ELISA.