

Human BST2 Protein

Cat. No. BST-HM202




Description	
Source	Recombinant Human BST2 Protein is expressed from HEK293 with hFc tag at the N-Terminus. It contains Asn49-Ser161.
Accession	Q10589-1
Molecular Weight	The protein has a predicted MW of 40 kDa. Due to glycosylation, the protein migrates to 50-60 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 24 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background	
Interferon-induced BST2 (bone marrow stromal cell antigen 2) inhibits viral replication by tethering enveloped virions to the cell surface to restrict viral release and by inducing the NFKB-dependent antiviral immune response. BST2 expression was significantly increased during porcine epidemic diarrhea virus (PEDV) infection of Vero cells by IRF1 targeting its promoter. Both the BST2 and N protein interacted with the E3 ubiquitin ligase MARCHF8/MARCH8 and the cargo receptor.	

Assay Data


Bis-Tris PAGE



Human BST2 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC


Assay Data



The purity of Human BST2 is greater than 95% as determined by SEC-HPLC.

ELISA Data

**Human BST2, hFc Tag ELISA**  
0.2μg Human BST2, hFc Tag Per Well



Immobilized Human BST2, hFc Tag at 2μg/ml (100μl/well) on the plate. Dose response curve for Biotinylated Anti-BST2 Antibody, hFc Tag with the EC50 of 40.0ng/ml determined by ELISA.