Human NCAM-1/CD56 Protein

Cat. No. CAM-HM256



Recombinant Human NCAM-1/CD56 Protein is expressed from HEK293 with hFc tag at the C-terminus.
It contains Leu20-Gly718.
P13591-2
The protein has a predicted MW of 103.33 kDa. Due to glycosylation, the protein migrates to 120-150 kDa based on Bis-Tris PAGE result.
Less than 1 EU per μg by the LAL method.
> 95% as determined by Bis-Tris PAGE
> 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22 μ m filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Neural Cell Adhesion Molecule 1 (NCAM-1), a multifunctional member of the immunoglobulin superfamily, is expressed on the surface of neurons, glia, skeletal muscle, and natural killer cells. NCAM-1 has been implicated as having a role in cell-cell adhesion, involved in development of the nervous system, and for cells involved in the expansion of T cells and dendritic cells which play an important role in immune surveillance.

Assay Data

Bis-Tris PAGE




Human NCAM-1 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC




Assay Data



The purity of Human NCAM-1 is greater than 95% as determined by SEC-HPLC.

ELISA Data

Human NCAM-1, hFc Tag ELISA 0.05μg Human NCAM-1, hFc Tag Per Well



Log Biotinylated Anti-NCAM-1 Antibody, hFc Tag Conc.(µg/ml)

Immobilized Human NCAM-1, hFc Tag at 0.5µg/ml (100µl/well) on the plate. Dose response curve for Biotinylated Anti-NCAM-1 Antibody, hFc Tag with the EC50 of 19.0ng/ml determined by ELISA.