

Human CD14 Protein

Cat. No. CD1-HM214




Description	
Source	Recombinant Human CD14 Protein is expressed from HEK293 with hFc tag at the C-terminus. It contains Thr20-Met344.
Accession	AAH10507
Molecular Weight	The protein has a predicted MW of 60.99 kDa. Due to glycosylation, the protein migrates to 70-80 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage	
Formulation	Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background	
Human monocyte differentiation antigen CD14 is a pattern recognition receptor (PRR) that enhances innate immune responses. CD14 was first identified as a marker of monocytes to signal intracellular responses upon bacterial encounters. Given the absence of an intracellular tail, CD14 was doubted to have the signaling capacities.	

Assay Data


Bis-Tris PAGE



Human CD14 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Assay Data



The purity of Human CD14 is greater than 95% as determined by SEC-HPLC.