

Human IL-2 R beta/CD122 Protein

Cat. No. CD1-HM422



Description

Source	Recombinant Human IL-2 R beta/CD122 Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus.
	It contains Ala27-Asp239.
Accession	NP_000869
Molecular Weight	The protein has a predicted MW of 27.5 kDa. Due to glycosylation, the protein migrates to 40-50 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

IL-2 R beta is a member of the cytokine receptor superfamily. Human IL-2 R beta cDNA encodes a 551 amino acid residue precursor Type I membrane protein with a 26 residue signal peptide, a 214 residue extracellular region, a 25 residue transmembrane region and a 286 residue cytoplasmic domain. Functional IL-2 receptors can exist in two affinity states on cell surfaces, the high affinity complex consisting of heterotrimers of the alpha, beta, and gamma chains, and the intermediate affinity complex comprising heterodimers of the beta and gamma chains.

Assay Data

Bis-Tris PAGE



Human IL-2 R beta on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.


SEC-HPLC

Human IL-2 R beta/CD122 Protein

Cat. No. CD1-HM422




Assay Data



The purity of Human IL-2 R beta is greater than 95% as determined by SEC-HPLC.


ELISA Data

Human IL-2 R beta, His Tag ELISA
0.2µg Human IL-2 R beta, His Tag Per Well



Immobilized Human IL-2 R beta, His Tag at 2µg/ml (100µl/well) on the plate. Dose response curve for Biotinylated Human IL-15Ra&IL-15, hFc Tag with the EC50 of 43.7ng/ml determined by ELISA (QC Test).

SPR Data



Human IL-2 R beta, His Tag captured on CM5 Chip via anti-his antibody can bind Human IL-2, No Tag with an affinity constant of 0.12 µM as determined in SPR assay (Biacore T200).


SPR Data

Human IL-2 R beta/CD122 Protein

Cat. No. CD1-HM422



Assay Data



Human IL-2 R beta, His Tag captured on CM5 Chip via Anti-His Antibody can bind Human IL-15, No Tag with an affinity constant of 5.9nM as determined in SPR assay (Biacore T200).