

Biotinylated Human CD24 Protein (Primary Amine Labeling)



Cat. No. CD2-HM324B


Description	
Source	Recombinant Biotinylated Human CD24 Protein (Primary Amine Labeling) is expressed from HEK293 with mFc (IgG1) tag at the C-Terminus. It contains Ser27-Gly59.
Accession	P25063-1
Molecular Weight	The protein has a predicted MW of 29.5 kDa. Due to glycosylation, the protein migrates to 48-58 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background	
CD24 is a sialoglycoprotein expressed at the surface of most B lymphocytes and differentiating neuroblasts. It is also expressed on neutrophils and neutrophil precursors from the myelocyte stage onwards. The potential for targeting CD24 in cancer therapy seems promising, as CD24 is overexpressed in many human cancers.	

Assay Data

Bis-Tris PAGE



Biotinylated Human CD24 on Bis-Tris PAGE under reduced conditions. The purity is greater than 95%.


SEC-HPLC

Biotinylated Human CD24 Protein (Primary Amine Labeling)

Cat. No. CD2-HM324B



Assay Data




The purity of Biotinylated Human CD24 is greater than 95% as determined by SEC-HPLC.

ELISA Data

Biotinylated Human CD24, mFc Tag ELISA

0.2µg Anti-CD24 Antibody, hFc Tag Per Well



Immobilized Anti-CD24 Antibody at 2µg/ml (100µl/Well) on the plate. Dose response curve for Biotinylated Human CD24, mFc Tag with the EC50 of 0.31µg/ml determined by ELISA.