

Cynomolgus CD3E&CD3D/CD3 epsilon&CD3 delta Protein



Cat. No. CD3-CM101


Description	
Source	Recombinant Cynomolgus CD3E&CD3D or CD3 epsilon&CD3 delta protein is expressed from HEK293 with His tag of CD3E and Flag tag of CD3D at the C-Terminus. It contains Gln22-Asp117 (CD3E) and Phe22-Ala105 (CD3D).
Accession	Q95LI5.2(CD3E)&Q95LI8(CD3D)
Molecular Weight	The protein has a predicted MW of 11.93 kDa (CD3E) and 10.55 kDa (CD3D). Due to glycosylation, the protein migrates to 13-20 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 0.1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE

Formulation and Storage	
Formulation	Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt.-80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background	
T-cell surface glycoprotein CD3 epsilon&CD3 delta chain, also known as CD3E&CD3D, are single-pass type I membrane proteins. When antigen presenting cells (APCs) activate T-cell receptor (TCR), TCR-mediated signals are transmitted across the cell membrane by the CD3 chains CD3D, CD3E, CD3G and CD3Z. All CD3 chains contain immunoreceptor tyrosine-based activation motifs (ITAMs) in their cytoplasmic domain.	

Assay Data


Bis-Tris PAGE



Cynomolgus CD3E&CD3D on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

ELISA Data

Cynomolgus CD3E&CD3D, His Tag ELISA
0.05µg Cynomolgus CD3E&CD3D, His Tag Per Well



Immobilized Cynomolgus CD3E&CD3D, His Tag at 0.5 µg/ml (100 µl/well) on the plate. Dose response curve for Anti-CD3 Antibody, hFc Tag with the EC50 of 5.6 ng/ml determined by ELISA (QC Test).