Mouse Siglec-3/CD33 Protein

Cat. No. CD3-MM133



| Description | |
|---------------------|---|
| Source | Recombinant Mouse Siglec-3/CD33 Protein is expressed from HEK293 with His tag at the C-Terminus. |
| | It contains Asp18-Glu240. |
| Accession | Q63994-1 |
| Molecular Weight | The protein has a predicted MW of 25.72 kDa. Due to glycosylation, the protein migrates to 40-50 kDa based on Bis-Tris PAGE result. |
| Endotoxin | Less than 1 EU per μg by the LAL method. |
| Purity | > 95% as determined by Bis-Tris PAGE |
| | > 95% as determined by HPLC |
| | |

Formulation and Storage


| Formulation | Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization. |
|----------------|---|
| Reconstitution | Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions. |
| Storage | -20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles. |

Background

Sialic-acid-binding immunoglobulin-like lectin (Siglec) that plays a role in mediating cell-cell interactions and in maintaining immune cells in a resting state. They are sialoadhesin/CD169/Siglec-1, CD22/Siglec-2, CD33/Siglec-3, Myelin-Associated Glycoprotein (MAG/Siglec-4a) and Siglecs 5 to 11. To date, no Siglec has been shown to recognized any cell surface ligand other than sialic acids, suggesting that interactions with glycans containing this carbohydrate are important in mediating the biological functions of Siglecs.

Assay Data

Bis-Tris PAGE




Mouse Siglec-3 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Cat. No. CD3-MM133



Assay Data



The purity of Mouse Siglec-3 is greater than 95% as determined by SEC-HPLC.